



Supplementary Materials

Catalytic performance of a new 1D Cu(II) coordination polymer {Cu(NO₃)(H₂O)}(HTae)(4,4'-Bpy) for Knoevenagel condensation

Edurne S. Larrea 1*, Roberto Fernández de Luis 2, and María I. Arriortua 1,2,

- Dpto. Mineralogía y Petrología, Universidad del País Vasco, UPV/EHU, Sarriena s/n, 48940 Leioa, Spain; edurne.serrano@ehu.eus, maribel.arriortua@ehu.eus
- ² BCMaterials (Basque Center for Materials, Applications & Nanostructures), Technological Park of Zamudio, Camino de Ibaizabal, Bndg. 500-1st, 48160, Derio, Spain.; roberto.fernandez@bcmaterials.net
- * Correspondence: edurne.serrano@ehu.eus; Tel.: +34-946-015-984




Figure S1. X-ray diffractograms of the activated **CuHTaeBpy_HT** catalyst before the reaction, after reacting with benzaldehyde (BA) and malononitrile (MN), and after reacting with benzaldehyde(BA) and ethyl cyanoacetate (ECA).








Figure S2. X-ray diffraction patterns of the activated catalyst, **CuHTaeBpy_HT**, and after reacting with various substrates and MN.




Figure S3. X-ray diffraction patterns of the activated catalyst, **CuHTaeBpy_HT**, and after 1 cycle of reaction and after 4 cycles of reaction of BA and MN.








Figure S4. IR spectra of the **CuHTaeBpy_RT** preactivated catalyst, the catalyst after reacting with benzaldehyde (BA) and malononitrile (MN), and after reacting with benzaldehyde (BA) and ethyl cyanoacetate (ECA).




Figure S5. IR spectra of the **CuHTaeBpy_RT** preactivated catalyst, the catalyst after reacting with various substrates and malononitrile.








Figure S6. Top: Rietveld refinements of the $\{Cu(NO_3)(H_2O)\}(HTae)(4,4'-Bpy)$ with the average structural model (left) and commensurate structural model (right). Bottom: Details of the refinement of the satellite peaks due to the super-structure.