

A SYSTEMATIC REVIEW AND META-ANALYSIS OF MTHFR POLYMORPHISMS IN METHOTREXATE TOXICITY PREDICTION IN PEDIATRIC ACUTE LYMPHOBLASTIC LEUKEMIA

Elixabet Lopez-Lopez, PhD (1); Idoia Martin-Guerrero, PhD (1); Javier Ballesteros, PhD (2); Africa Garcia-Orad, PhD (1)

(1)Department of Genetics, Physic Anthropology and Animal Physiology, University of the Basque Country, Leioa, Spain.

(2)Department of Neurosciences, University of the Basque Country, Leioa, CIBERSAM, Spain.

Correspondence

Africa Garcia-Orad, Department of Genetics, Physic Anthropology and Animal Physiology, Faculty of Medicine and Dentistry-University of the Basque Country, Barrio Sarriena s/n, 48940 Leioa, Spain.

Phone: international +34.946012909. Fax: international +34.946013400

E-mail: africa.garciaorad@ehu.es

Running title

MTHFR SNPS IN MTX TOXICITY IN PEDIATRIC ALL

Acknowledgments

This project was supported by RTICC (RD/06/0020/0048), Basque Government (GIC10/71, SAI10/03), and UPV/EHU (UFI11/35). ELL was supported by a predoctoral grant from the Basque Government.

Conflict of Interest Statement

The authors reported no potential conflicts of interest.

Abstract

Methotrexate (MTX) is an important component of therapy used to treat childhood acute lymphoblastic leukemia (ALL). Two single nucleotide polymorphisms (SNPs) in the methylenetetrahydrofolate (*MTHFR*) gene, C677T and A1298C, affect MTHFR activity. A large body of studies has investigated the potential role of *MTHFR* SNPs in MTX toxicity in pediatric ALL. However, the results are controversial.

In this review and meta-analysis we critically evaluate the relationship between the C677T and A1298C polymorphisms of *MTHFR* and MTX toxicity in pediatric ALL.

The majority of published reports do not find associations between *MTHFR* polymorphisms and toxicity in pediatric ALL. When associations are reported, often the results are contradictory to each other. The meta-analysis confirms a lack of association. In conclusion, *MTHFR* C677T and A1298C polymorphisms do not seem to be good markers of MTX-related toxicity in pediatric ALL.

Keywords

Pharmacogenetics, polymorphism, methotrexate, MTHFR, acute lymphoblastic leukemia, toxicity

Introduction

Acute lymphoblastic leukemia (ALL) is the most common childhood cancer, accounting for 30% of all pediatric malignancies^{1,2}. During the last 20 years, survival rates for ALL have improved dramatically due to advances in specific chemotherapy for childhood ALL, with expected cure rates higher than 80%³.

Methotrexate (MTX) is an important component of the therapy for childhood ALL. Despite clinical success, treatment with MTX often causes toxicity, requiring a dose reduction or cessation of treatment. An accurate predictor of the adverse effects of MTX treatment would therefore be very useful in ALL⁴.

MTX enters the cell via active transport mediated by the reduced folate carrier (RFC1)⁵. Then, MTX acts by inhibiting mainly two enzymes. MTX inhibits dihydrofolate reductase (DHFR), inhibiting the folic acid cycle and affecting other important enzymes, such as methylenetetrahydrofolate reductase (MTHFR) and serine hydromethyl transferase (SHMT1). On the other hand, the conversion of MTX to its polyglutamated forms (MTXPG) results in the inhibition of thymidylate synthase (TS). These combined mechanisms interfere with nucleic acid synthesis, favouring cell death⁶. Effective cellular levels of MTX are reduced by different transporters that pump MTX out of the organism. These transporters include ABC transporters, such as the multidrug resistance protein (ABCB1) and the breast cancer resistance protein (ABCG2)⁷, and organic anion transporters, such as SLCO1B1^{8,9} (Figure 1).

MTHFR is a key enzyme for intracellular folate homeostasis and metabolism. MTHFR catalyses the irreversible conversion of 5,10-methylenetetrahydrofolate (5,10-CH₂-THF), required for

purine and thymidine synthesis, to 5-methyltetrahydrofolate (5-CH-THF), which is required for protein synthesis and nucleic acid methylation. Alterations in reduced folate pools, as a consequence of changes in MTHFR activity, may have a significant effect on the responsiveness of malignant and non-malignant cells to MTX. Accordingly, it has been proposed that an impaired conversion of 5,10-CH₂-THF to 5-CH-THF, and the subsequent modification in the intracellular folates pool could increase the toxic effect of MTX ¹⁰.

In this context, the non-synonymous SNPs C677T (causing Ala222Val) and A1298C (causing Glu429Ala) in the *MTHFR* gene have been widely studied. The *MTHFR* 677T allele encodes proteins with decreased enzymatic activity, in comparison with the normal allele 677C. People with the 677CT and 677TT genotype exhibit 60% and 30%, respectively, of the normal MTHFR activity ^{11, 12}. In the *MTHFR* A1298C polymorphism, the 1298C allele is responsible for a milder decrease in MTHFR activity with respect to the normal allele 1298A. The 1298CC homozygous individuals have 60% of the normal activity ¹³.

A large body of published studies has investigated the potential role of *MTHFR* polymorphisms in susceptibility, toxicity and response to MTX in pediatric ALL, with conflicting results. Possible reasons for these discrepancies are differences in treatment protocols among studies, small or non-homogeneous populations, ethnic differences, and the use of different criteria defining toxicity.

In this study, we have performed a critical review of the published articles on the relationship between genetic variants of *MTHFR* and the toxicity of MTX in pediatric ALL. Then, we undertook a meta-analysis on all eligible studies, separating them by toxicity criteria, to determine the role of the *MTHFR* C677T and A1298C polymorphisms on MTX toxicity in this pediatric ALL patients.

Methods

Search strategy

We performed an exhaustive search to identify studies that examined the association between the C677T and A1298C polymorphisms of *MTHFR* and MTX toxicity in pediatric ALL patients. We used the keywords and subject terms “MTHFR and acute leukemia”, and “MTHFR and polymorphism(s) and toxicity” to search Pubmed (www.ncbi.nlm.nih.gov/pubmed) for articles published through November 2011. All references within the identified studies were then reviewed to possibly identify additional works.

Meta-analysis

Inclusion and exclusion criteria

The inclusion criteria for meta-analysis required that each trial be an independent association study, that the article supplied enough information on toxicity by genotype, that it studied short term toxic effects and that the population was composed only of pediatric patients (< 18 y). An article was excluded from meta-analysis if the study did not provide enough information (incomplete summary data), was performed on adult patients, the diagnosis was not mainly ALL or was a case study.

Data Extraction

For each article included in the study, we gathered ethnicity of study population, patients number, age and diagnosis, MTX dose, *MTHFR* C677T and A1298C genotype data and toxicity types.

Statistical analysis

Statistical analysis was performed using R software using the meta library (R version 2.11.0. The R Foundation for Statistical Computing). We used a recessive model, assuming a recessive effect of the minor allele of each *MTHFR* SNP, which was consistent with previous results and allowed inclusion of the maximum number of studies. For the C677T SNP, we compared individuals having the TT homozygous genotype to all others (CC + CT), and for the A1298C SNP we compared CC homozygous individuals to all others (AA + AC). The overall pooled relative risk (RR) and corresponding 95% CI of toxicity to MTX were estimated using Mantel–Haenszel’s method with random effect model. The random effects model assumes different underlying effects, considering both within- and between-study variation, offering an advantage as it accommodates diversity between studies and provides a more conservative estimate of the assessed effect.

Heterogeneity of the studies was assessed using the Cochran Q test with a P-value below 0.05, below which heterogeneity was considered statistically significant. The heterogeneity was also quantified using the I^2 statistic, which is independent of the number of studies in the meta-analysis. This statistic quantifies the effect of heterogeneity, providing a measure of the degree of inconsistency in the study’s results. The I^2 statistic has a value between 0 and 100% and describes the percentage of total variation across studies that is due to between-studies heterogeneity rather than chance. A higher I^2 value denotes a greater degree of heterogeneity (customary interpretations of the I^2 value are, 0–25% no heterogeneity, 25–50% moderate heterogeneity, 50–75% large heterogeneity, 75–100% extreme heterogeneity). Sensitivity analysis leaving out one study at the time was also performed when possible: outlying studies were identified and excluded and the I^2 estimates for these different sets of studies examined.

Results

The original search provided 264 records. After eliminating duplications, 238 records remained. Of these, 117 were discarded after reviewing the abstracts because they clearly did not meet the required criteria for inclusion. The full texts of the remaining 121 studies were examined in detail. Of these, we identified 24 studies which investigated *MTHFR* SNPs and MTX related toxicity in pediatric ALL patients for meta-analysis (Figure 2). All 24 studied the C677T polymorphism (Table 1) and 16 of these also studied the A1298C polymorphism (Table 2).

In general, the 24 studies could be categorized according to the level of association between *MTHFR* SNPs and MTX toxicity: those that found no association, those that found an association between *MTHFR* SNP and a significant increase in toxicity, and those that found an association between *MTHFR* SNP and a significant decrease in toxicity. No population, even grouped by ethnicity or geography, was overrepresented in any of the 3 groups of studies. Additionally, both high and low dose MTX were found in all three studied groups. We did not find either differences between the reports that used normal or tumor samples (Tables 1-2). Because different studies analyzed toxicity according to different criteria, we performed in-depth analysis for each toxicity criterion (Tables 3, 4).

MTHFR C677T polymorphism and toxicity in pediatric ALL

We could not perform a meta-analysis for treatment interruption, MTX clearance, diarrhea, hyperbilirubinemia and renal toxicity due to lack of data. We did perform meta-analysis for MTX plasma levels, mucositis, hepatic toxicity, neutropenia, thrombocytopenia, anemia and leucopenia.

MTX plasma levels:

MTX plasma levels were studied in five works that used high MTX doses ^{4, 14-17}. Only 2 studies provided enough data, from a total of 137 patients, to be included in the meta-analysis ^{4, 15}. We found no statistical association between C677T and MTX plasma levels (Figure 3).

Mucositis:

Mucositis was surveyed in 10 studies ^{14, 17-25}. One of them was performed with low MTX doses ²¹ and the rest with high doses. We performed meta-analysis on 4 studies ^{17, 19, 23, 26} with data from a total of 484 observations. No association with mucositis was observed (Figure 3). As the heterogeneity among studies was high, a sensitivity analysis was undertaken and this identified the study by Tantawy et al. ²⁶ as an outlier. Removing this data from the meta-analysis reduced the heterogeneity, yet the pooled RR remained non significant.

Hepatic toxicity:

We compiled 16 studies that analyzed hepatic toxicity with low ^{21, 27, 28} or high MTX doses ^{4, 14, 16-19, 22, 25, 26, 29-32}. Of these 16 studies, 6 presented enough data to allow meta-analysis ^{4, 17, 19, 26, 27, 31} with data from a total of 757 patients. No association between C677T genotypes and hepatic toxicity was observed (Figure 3). Since there was a great heterogeneity between studies, a sensitivity analysis was undertaken and this identified the study by Tantawy et al. ²⁶ as an outlier. Removing this data from the meta-analysis reduced the heterogeneity yet the pooled RR remained non significant. We also performed a meta-analysis with the 5 high dose studies that provided data and we obtained similar results.

Neutropenia:

From the 4 papers that analyzed neutropenia with low ²⁸ or high MTX doses ^{17, 26, 30} (Table 1), 2 provided enough data to be included in the meta-analysis ^{17, 26} with data from 200 patients. No association between C677T SNP and neutropenia was observed (Figure 3).

Thrombocytopenia:

A total of 10 studies^{14, 16, 17, 19, 22, 23, 25, 28, 30, 32} analyzed thrombocytopenia. One of them was performed with low MTX doses²⁸ and the rest with high doses. In the meta-analysis, 3 studies were included^{17, 19, 23} with data from a total of 381 observations. No association between the C677T SNP and thrombocytopenia was observed (Figure 3).

Anemia:

From the 8 reports that studied anemia^{14, 16, 17, 19, 22, 25, 29, 32} with high MTX doses, a single study³² found an association between C677T and increased anemia. In the meta-analysis, we excluded 5 studies due to lack of data, leaving 2 studies with data from 192 patients^{17, 19}. We observed no association with anemia (Figure 3).

Leucopenia:

We found 10 reports that studied leucopenia with low²⁸ or high MTX doses^{14, 16, 19, 22, 23, 25, 29, 30, 32}. From these 10 studies, 2 provided genotype data from 221 observations^{19, 23}. No association between C677T and leucopenia in response to MTX treatment in ALL was observed (Figure 3).

MTHFR A1298C polymorphism and toxicity in pediatric ALL

In the 16 studies that analyzed this polymorphism (Table 2), 8 studies with low^{21, 28, 33} or high MTX doses^{15, 22, 26, 34, 35} found no association between A1298C and any toxic effect. In 5 studies, with high MTX doses, the authors reported a protective effect of the 1298C allele against various types of MTX toxicity^{14, 17, 20, 23, 30}. We found three studies, also with high MTX doses, in which this allele was associated with higher MTX toxicity^{16, 19, 25}.

We could not perform a meta-analysis for transfusions, skin toxicity, MTX plasma levels, or febrile neutropenia due to lack of data. We did perform meta-analyses for leucopenia, myelosuppression, thrombocytopenia, hepatic toxicity, and anemia with high MTX doses and only observed a slight protective effect of the 1298CC genotype for leucopenia in a meta-analysis study with data from only two reports (Figure 4).

Discussion

MTHFR C677T polymorphism and toxicity in pediatric ALL

In the 24 published studies used in this analysis, 12 did not find a significant association between the *MTHFR* 677T low functional allele and MTX toxicity^{14, 15, 18-22, 27, 29, 33, 34, 36}. Three studies found an association between the 677T allele and a decrease in toxicity^{16, 28, 30}. Nine studies found an association between this allele and increased toxicity^{4, 17, 23, 25, 26, 31, 32, 35, 37} (Table 1).

Populations studied or MTX doses could not explain the differences in results among studies. We did not find either differences between the reports that used normal or tumor samples, as it was expected considering that mutations or deletions have not been described in *MTHFR* in ALL. Because different studies analyzed toxicity according to different criteria, below we analyze the findings from the 24 studies for each toxicity criterion and report results from meta-analysis if enough data was provided to make it possible.

Treatment interruption:

Three studies analyzed MTX treatment interruption. An association between the 677T allele and an increase in interruption was reported by Shimasaki et al with low MTX doses³⁷, however this study was carried out with a small and heterogeneous population (20 ALL or lymphoblastic lymphoma (LBL)) and only one patient with the TT genotype was reported. Two

larger studies of 88 and 201 ALL patients did not find any association between 677T and MTX treatment interruption with low³³ or high MTX doses³⁰. Consequently, the 677TT genotype cannot be considered a good predictor of treatment interruption. The three articles did not provide enough information to carry out a meta-analysis to confirm it.

MTX pharmacokinetics:

Although MTX pharmacokinetics are more directly related with membrane transporters, they have been measured in association with MTHFR C677T polymorphism in several studies using high MTX doses with different parameters: MTX plasma levels, MTX clearance and renal toxicity.

MTX plasma levels were studied in five works. Imanishi et al studied 26 children with ALL or malignant lymphoma (ML)⁴ and concluded that patients with the 677TT homozygous genotype had higher MTX plasma levels 48 h after infusion. The other 4 studies found no association between the C677T SNP and MTX plasma levels 48 h or 72 h after infusion¹⁴⁻¹⁷. We did not find either any statistical association between C677T and MTX plasma levels in the meta-analysis.

Two studies analyzed MTX clearance and reported conflicting results. One study of 64 children with LLA or ML²³ found an association between the 677TT homozygous genotype and a decrease in MTX clearance. The larger study of 240 pediatric ALL patients did not find any association between the C677T SNP and MTX clearance³⁴. We could not carry out a meta-analysis for this parameter.

Renal toxicity was reported in 3 studies. One³² found an association between the 677T allele and increased renal toxicity. In a combined study of C677T and A1298C, association with

increased renal toxicity was also found ²⁵. Another larger study did not find any association between the 677T allele and increased renal toxicity ²². Consequently, the published data do not support a clear association between the 677T allele and renal toxicity in response to MTX treatment in ALL. We could not carry out a meta-analysis to confirm it, due to lack of data.

Consequently, the 677T allele does not seem to be a good marker of MTX pharmacokinetics in ALL treatment.

Gastrointestinal toxicity:

Some studies found associations between the *MTHFR* 677T allele and gastrointestinal toxicity parameters: diarrhea and mucositis.

Four studies analyzed diarrhea with high MTX doses. An association between the 677TT homozygous genotype and higher risk of diarrhea was found in a single study of 40 pediatric ALL patients ²⁶. Three additional studies carried out with 240, 520, and 557 pediatric ALL patients did not find this correlation ^{21, 31, 34}. Accordingly, the 677TT genotype cannot be considered a good predictor of severe diarrhea in response to MTX treatment for ALL. Only one of the 4 articles provided genotype information, so we were unable to confirm this with a meta-analysis.

Mucositis was surveyed in 10 studies. Two studies of 40 and 64 children with ALL found an association between 677TT genotype and higher risk of mucositis with high MTX doses ^{23, 26}. The other 8 studies, most of which were larger, studied various ethnic populations and included low ²¹ and high MTX doses ^{14, 17-20, 22, 25}, did not find this association. This lack of consistent results across these studies does not support an effect of the 677TT genotype in the

risk of mucositis in response to MTX treatment for ALL. Our meta-analysis supported this lack of association.

Hepatic toxicity:

We compiled 16 studies that analyzed hepatic toxicity (transaminitis). Three of them found an association between the 677TT genotype and increased hepatic toxicity with high MTX doses^{26, 31, 32}. However, two of these studies do not have a very high statistical power, and the other 13 studies that analyzed this parameter found no association between 677TT genotype and hepatic toxicity with low^{21, 27, 28} or high MTX doses^{4, 14, 16-19, 22, 25, 29, 30}, therefore we conclude that the 677TT genotype does not appear to be a good predictor of hepatic toxicity in response to MTX treatment for ALL. This lack of association was confirmed with the meta-analyses performed.

Hyperbilirubinemia, which is also associated with hepatic toxicity, was studied in four reports. One study of 37 patients¹⁶ found that individuals with 677CT or 677TT genotypes had less hyperbilirubinemia with high MTX doses. Three larger studies of 240, 520 and 557 patients did not find this association with low²¹ or high MTX doses^{31, 34}. None of these articles provided enough information to perform a meta-analysis. We conclude that the 677TT genotype does not appear to be either a good predictor of hyperbilirubinemia in response to MTX treatment for ALL.

Hematologic toxicity:

Some studies have found associations between *MTHFR* 677T allele and different hematologic toxicity parameters: neutropenia, thrombocytopenia, anemia and leucopenia.

From the 4 papers that analyzed neutropenia, only one reported an association between the 677TT genotype and higher risk of neutropenia with high MTX doses ²⁶. Two larger studies did not find this association with low ²⁸ or high MTX doses ¹⁷. A fourth study reported the opposite effect, finding an association between the 677TT genotype and a lower risk of neutropenia ³⁰ with high MTX doses (Table 1). The controversial reports and the results of our meta-analysis do not support an association between C677T SNP and neutropenia.

A total of 10 studies analyzed thrombocytopenia with low ²⁸ or high MTX doses ^{14, 16, 17, 19, 22, 23, 25, 30, 32}. An association between the 677CT and 677TT genotypes and an increased risk of thrombocytopenia was reported in 2 studies ^{17, 32}, but was only statistically significant for the 677CT genotype. The apparent disadvantage of the heterozygous genotype is difficult to explain from a functional point of view. Furthermore, another study reported a correlation between the 677CT and TT genotypes with decreased risk of thrombocytopenia ¹⁶. In another study that looked at both C677T and A1298C, an association between the combined 677T and 1298C alleles and increased thrombocytopenia was found ²⁵. An additional 6 studies did not find any association between C677T SNP and thrombocytopenia ^{14, 19, 22, 23, 28, 30}. In conclusion, the available data do not support a clear association between the 677T allele and a higher risk of thrombocytopenia in response to MTX treatment for ALL, which is in agreement with our meta-analysis.

From the 8 reports that studied anemia with high MTX doses ^{14, 16, 17, 19, 22, 25, 29, 32}, a single study ³² found an association between C677T and increased anemia. In the meta-analysis, we observed no association with anemia.

We also found 10 reports that studied leucopenia. One ³² found an association between C677T and increased leucopenia with high MTX doses. One study reported the opposite, finding an

association between 677T and decreased leucopenia with low MTX doses ²⁸. 8 studies did not find any association with high MTX doses ^{14, 16, 19, 22, 23, 25, 29, 30}. These results and our meta-analysis do not support a clear association between 677T allele and leucopenia.

Consequently, MTHFR 677T allele does not seem to be a good marker of hematologic toxicity.

Finally, in our review of the literature, we re-analyzed, when possible, the data provided in the articles. In one case in which the authors reported an association between the 677TT genotype and an increase in global toxicity ³⁵, we detected a statistical error and drew the opposite conclusion to what the authors proposed ³⁸.

MTHFR A1298C polymorphism and toxicity in pediatric ALL

In the 16 studies that analyzed this polymorphism (Table 2), 8 studies ^{15, 21, 22, 26, 28, 33-35} found no association between A1298C and any toxic effect. In 5 studies, the authors reported a protective effect of the 1298C allele against various types of MTX toxicity ^{14, 17, 20, 23, 30}. We found three studies in which this allele was associated with higher MTX toxicity ^{16, 19, 25}. In our meta-analyses, we only observed a slight protective effect of the 1298CC genotype for leucopenia in a meta-analysis study with data from only two reports.

According to the published data and the meta-analysis reported, the 1298C allele does not seem to be a good MTX toxicity marker in pediatric ALL patients. If anything, it seems to be more likely a protective factor rather than a toxicity marker.

In summary, considering all the previously discussed combined with the fact that works that could not be included in the meta-analyses due to lack of data, are, in general, those which

found no association with toxicity, we conclude that there is no evidence to support the use of either the *MTHFR* C677T or the A1298C SNP as MTX toxicity markers.

Taking into account our conclusion that *MTHFR* C677T and A1298C SNPs do not seem good MTX toxicity markers and that a recent study reported that patients receiving higher doses of MTX have better survival ²⁵, patients with pediatric ALL might benefit from higher MTX doses in spite of their *MTHFR* genotype.

Acknowledgments

This project was supported by RTICC (RD/06/0020/0048), Basque Government (GIC10/71, SAI10/03), and UPV/EHU (UFI11/35). ELL was supported by a predoctoral grant from the Basque Government.

Conflict of Interest Statement

The authors reported no potential conflicts of interest.

References

1. Koppen IJ, Hermans FJ, Kaspers GJ. Folate related gene polymorphisms and susceptibility to develop childhood acute lymphoblastic leukaemia. *Br J Haematol* 2010; **148**(1): 3-14.
2. Johnston WT, Lightfoot TJ, Simpson J, Roman E. Childhood cancer survival: a report from the United Kingdom Childhood Cancer Study. *Cancer Epidemiol* 2010; **34**(6): 659-666.
3. Pui CH, Robison LL, Look AT. Acute lymphoblastic leukaemia. *Lancet* 2008; **371**(9617): 1030-1043.
4. Imanishi H, Okamura N, Yagi M, Noro Y, Moriya Y, Nakamura T, *et al.* Genetic polymorphisms associated with adverse events and elimination of methotrexate in childhood acute lymphoblastic leukemia and malignant lymphoma. *J Hum Genet* 2007; **52**(2): 166-171.
5. Gorlick R, Goker E, Trippett T, Waltham M, Banerjee D, Bertino JR. Intrinsic and acquired resistance to methotrexate in acute leukemia. *N Engl J Med* 1996; **335**(14): 1041-1048.
6. Krajcinovic M, Moghrabi A. Pharmacogenetics of methotrexate. *Pharmacogenomics* 2004; **5**(7): 819-834.

7. Swerts K, De Moerloose B, Dhooge C, Laureys G, Benoit Y, Philippé J. Prognostic significance of multidrug resistance-related proteins in childhood acute lymphoblastic leukaemia. *Eur J Cancer* 2006; **42**(3): 295-309.
8. Treviño LR, Shimasaki N, Yang W, Panetta JC, Cheng C, Pei D, *et al.* Germline genetic variation in an organic anion transporter polypeptide associated with methotrexate pharmacokinetics and clinical effects. *J Clin Oncol* 2009; **27**(35): 5972-5978.
9. Abe T, Unno M, Onogawa T, Tokui T, Kondo TN, Nakagomi R, *et al.* LST-2, a human liver-specific organic anion transporter, determines methotrexate sensitivity in gastrointestinal cancers. *Gastroenterology* 2001; **120**(7): 1689-1699.
10. De Mattia E, Toffoli G. C677T and A1298C MTHFR polymorphisms, a challenge for antifolate and fluoropyrimidine-based therapy personalisation. *Eur J Cancer* 2009; **45**(8): 1333-1351.
11. Cheok MH, Evans WE. Acute lymphoblastic leukaemia: a model for the pharmacogenomics of cancer therapy. *Nat Rev Cancer* 2006; **6**(2): 117-129.
12. Frosst P, Blom HJ, Milos R, Goyette P, Sheppard CA, Matthews RG, *et al.* A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nat Genet* 1995; **10**(1): 111-113.

13. Weisberg IS, Jacques PF, Selhub J, Bostom AG, Chen Z, Curtis Ellison R, *et al.* The 1298A-->C polymorphism in methylenetetrahydrofolate reductase (MTHFR): in vitro expression and association with homocysteine. *Atherosclerosis* 2001; **156**(2): 409-415.
14. Huang L, Tissing WJ, de Jonge R, van Zelst BD, Pieters R. Polymorphisms in folate-related genes: association with side effects of high-dose methotrexate in childhood acute lymphoblastic leukemia. *Leukemia* 2008; **22**(9): 1798-1800.
15. Lopez-Lopez E, Martin-Guerrero I, Ballesteros J, Piñan MA, Garcia-Miguel P, Navajas A, *et al.* Polymorphisms of the SLCO1B1 gene predict methotrexate-related toxicity in childhood acute lymphoblastic leukemia. *Pediatr Blood Cancer* 2011; **57**(4): 612-619.
16. Kantar M, Kosova B, Cetingul N, Gumus S, Toroslu E, Zafer N, *et al.* Methylenetetrahydrofolate reductase C677T and A1298C gene polymorphisms and therapy-related toxicity in children treated for acute lymphoblastic leukemia and non-Hodgkin lymphoma. *Leuk Lymphoma* 2009; **50**(6): 912-917.
17. Liu SG, Li ZG, Cui L, Gao C, Li WJ, Zhao XX. Effects of methylenetetrahydrofolate reductase gene polymorphisms on toxicities during consolidation therapy in pediatric acute lymphoblastic leukemia in a Chinese population. *Leuk Lymphoma* 2011; **52**(6): 1030-1040.
18. Shimasaki N, Mori T, Samejima H, Sato R, Shimada H, Yahagi N, *et al.* Effects of methylenetetrahydrofolate reductase and reduced folate carrier 1 polymorphisms on

- high-dose methotrexate-induced toxicities in children with acute lymphoblastic leukemia or lymphoma. *J Pediatr Hematol Oncol* 2006; **28**(2): 64-68.
19. Karathanasis NV, Stiakaki E, Goulielmos GN, Kalmanti M. The role of the methylenetetrahydrofolate reductase 677 and 1298 polymorphisms in Cretan children with acute lymphoblastic leukemia. *Genet Test Mol Biomarkers* 2011; **15**(1-2): 5-10.
 20. Pakakasama S, Kanchanakamhaeng K, Kajanachumpol S, Udomsubpayakul U, Sirachainan N, Thithapandha A, *et al.* Genetic polymorphisms of folate metabolic enzymes and toxicities of high dose methotrexate in children with acute lymphoblastic leukemia. *Ann Hematol* 2007; **86**(8): 609-611.
 21. Aplenc R, Thompson J, Han P, La M, Zhao H, Lange B, *et al.* Methylenetetrahydrofolate reductase polymorphisms and therapy response in pediatric acute lymphoblastic leukemia. *Cancer Res* 2005; **65**(6): 2482-2487.
 22. Erčulj N, Kotnik BF, Debeljak M, Jazbec J, Dolžan V. Influence of folate pathway polymorphisms on high-dose methotrexate-related toxicity and survival in childhood acute lymphoblastic leukemia. *Leuk Lymphoma* 2012.
 23. Faganel Kotnik B, Grabnar I, Bohanec Grabar P, Dolžan V, Jazbec J. Association of genetic polymorphism in the folate metabolic pathway with methotrexate pharmacokinetics and toxicity in childhood acute lymphoblastic leukaemia and malignant lymphoma. *Eur J Clin Pharmacol* 2011; **67**(10): 993-1006.

24. Faganel Kotnik B, Dolzan V, Grabnar I, Jazbec J. Relationship of the reduced folate carrier gene polymorphism G80A to methotrexate plasma concentration, toxicity, and disease outcome in childhood acute lymphoblastic leukemia. *Leuk Lymphoma* 2010; **51**(4): 724-726.
25. Salazar J, Altés A, Del Río E, Estella J, Rives S, Tasso M, *et al.* Methotrexate consolidation treatment according to pharmacogenetics of MTHFR ameliorates event-free survival in childhood acute lymphoblastic leukaemia. *Pharmacogenomics J* 2011.
26. Tantawy AA, El-Bostany EA, Adly AA, Abou El Asrar M, El-Ghouroury EA, Abdulghaffar EE. Methylene tetrahydrofolate reductase gene polymorphism in Egyptian children with acute lymphoblastic leukemia. *Blood Coagul Fibrinolysis* 2010; **21**(1): 28-34.
27. Horinouchi M, Yagi M, Imanishi H, Mori T, Yanai T, Hayakawa A, *et al.* Association of genetic polymorphisms with hepatotoxicity in patients with childhood acute lymphoblastic leukemia or lymphoma. *Pediatr Hematol Oncol* 2010; **27**(5): 344-354.
28. Costea I, Moghrabi A, Laverdiere C, Graziani A, Krajinovic M. Folate cycle gene variants and chemotherapy toxicity in pediatric patients with acute lymphoblastic leukemia. *Haematologica* 2006; **91**(8): 1113-1116.
29. Chatzidakis K, Goulas A, Athanassiadou-Piperopoulou F, Fidani L, Kolioukas D, Mirtsou V. Methylene tetrahydrofolate reductase C677T polymorphism: association with risk

- for childhood acute lymphoblastic leukemia and response during the initial phase of chemotherapy in greek patients. *Pediatr Blood Cancer* 2006; **47**(2): 147-151.
30. van Kooten Niekerk PB, Schmiegelow K, Schroeder H. Influence of methylene tetrahydrofolate reductase polymorphisms and coadministration of antimetabolites on toxicity after high dose methotrexate. *Eur J Haematol* 2008; **81**(5): 391-398.
 31. Sepe DM, McWilliams T, Chen J, Kershenbaum A, Zhao H, La M, *et al.* Germline genetic variation and treatment response on CCG-1891. *Pediatr Blood Cancer* 2012; **58**(5): 695-700.
 32. El-Khodary NM, El-Haggar SM, Eid MA, Ebeid EN. Study of the pharmacokinetic and pharmacogenetic contribution to the toxicity of high-dose methotrexate in children with acute lymphoblastic leukemia. *Med Oncol* 2011.
 33. Krajcinovic M, Lemieux-Blanchard E, Chiasson S, Primeau M, Costea I, Moghrabi A. Role of polymorphisms in MTHFR and MTHFD1 genes in the outcome of childhood acute lymphoblastic leukemia. *Pharmacogenomics J* 2004; **4**(1): 66-72.
 34. Kishi S, Cheng C, French D, Pei D, Das S, Cook EH, *et al.* Ancestry and pharmacogenetics of antileukemic drug toxicity. *Blood* 2007; **109**(10): 4151-4157.
 35. D'Angelo V, Ramaglia M, Iannotta A, Crisci S, Indolfi P, Francese M, *et al.* Methotrexate toxicity and efficacy during the consolidation phase in paediatric acute lymphoblastic

leukaemia and MTHFR polymorphisms as pharmacogenetic determinants. *Cancer Chemother Pharmacol* 2011; **68**(5): 1339-1346.

36. Kishi S, Griener J, Cheng C, Das S, Cook EH, Pei D, *et al.* Homocysteine, pharmacogenetics, and neurotoxicity in children with leukemia. *J Clin Oncol* 2003; **21**(16): 3084-3091.

37. Shimasaki N, Mori T, Torii C, Sato R, Shimada H, Tanigawara Y, *et al.* Influence of MTHFR and RFC1 polymorphisms on toxicities during maintenance chemotherapy for childhood acute lymphoblastic leukemia or lymphoma. *J Pediatr Hematol Oncol* 2008; **30**(5): 347-352.

38. Lopez-Lopez E, Ballesteros J, Garcia-Orad A. MTHFR 677TT genotype and toxicity of methotrexate: controversial results. *Cancer Chemother Pharmacol* 2011; **68**(5): 1369-1370; author reply 1371.

Figure 1. Methotrexate pathway. Methotrexate and its metabolites are indicated. Enzymes and transporters are encircled. *MTHFR* is in black.

Figure 2. Flow diagram of study selection.

Figure 3. Results of meta-analysis of association between the *MTHFR* C677T SNP and MTX toxicities in treatment of ALL. No associations were confirmed between genotype and toxicity.

Figure 4. Results of meta-analysis of association between the *MTHFR* A1298C polymorphism and MTX toxicities in treatment of ALL. We observed a slight protective effect of the 1298CC genotype with leucopenia using data from only two reports.

Table 1. List of 24 studies that analyzed association between the *MTHFR* C677T polymorphism and MTX toxicity in pediatric ALL, grouped according to the level of association between the SNP and MTX toxicity

MTHFR C677T				
Patient population	MTX dose	Population	Association with toxicity	Reference
15 ALL or LBL ^A	high	Japanese	NA	Shimasaki et al, 2006 ¹⁸
24 ALL or LBL ^A	low	Japanese	NA	Horinouchi et al, 2010 ²⁷
35 ALL ^C	high	Cretan	NA	Karathanasis et al, 2011 ¹⁹
46 ALL ^C	high	Greek	NA	Chatzidakis et al, 2006 ²⁹
53 ALL ^C	high	Various	NA	Kishi et al, 2003 ³⁶
76 ALL ^C	high	Thai	NA	Pakakasama et al, 2007 ²⁰
81 ALL ^B	high	European	NA	Huang et al, 2008 ¹⁴
115 ALL ^A	high	Spanish	NA	Lopez-Lopez et al, 2011 ¹⁵
167 ALL ^C	high	European	NA	Erculj et al, 2012 ²²
240 ALL ^A	high	North American	NA	Kishi et al, 2007 ³⁴
201 ALL ^C	low	French-Canadian	NA	Krajinovic et al, 2004 ³³
520 ALL ^B	low	Various	NA	Aplenc et al, 2005 ²¹
37 ALL or NHL ^C	high	Turkish	-T	Kantar et al, 2009 ¹⁶
88 ALL ^C	high	European	-T	van Kooten et al, 2008 ³⁰
186 ALL ^B	low	European	-T	Costea et al, 2006 ²⁸
20 ALL or LBL ^A	low	Japanese	+T	Shimasaki et al, 2008 ³⁷
26 ALL or ML ^A	high	Japanese	+T	Imanishi et al, 2007 ⁴
40 ALL ^C	high	Egyptian	+T	Tantawy et al, 2010 ²⁶
40 ALL ^A	high	Egyptian	+T	EL-Khodary et al, 2011 ³²
64 ALL or ML ^A	high	European	+T	Faganel Kotnik et al, 2011 ²³
141 ALL ^C	high	Spanish	+T	Salazar et al, 2011 ²⁵
151 ALL ^C	high	European	+T	D'Angelo et al, 2011 ³⁵
181 ALL ^B	high	Chinese	+T	Liu et al, 2011 ¹⁷
557 ALL ^C	high	Various	+T	Sepe et al, 2012 ³¹

Type of sample: A: normal, B: tumor, C: unknown

High MTX dose = 1.5 – 5 g / m²; Low MTX dose = 15 – 30 mg / m²

NA, no association between the SNP and toxicity

+T, SNP is associated with increased toxicity (light shading)

-T, SNP is associated with decreased toxicity (dark shading)

Table 2. Association of *MTHFR* A1298C polymorphism and toxicity in pediatric acute lymphoblastic leukemia

MTHFR A1298C				
Patient population	MTX dose	Population	Association with toxicity	Reference
40 LLAs ^C	high	Egyptian	NA	Tantawy et al, 2010 ²⁶
115 LLA ^A	high	Spanish	NA	Lopez-Lopez et al, 2011 ¹⁵
151 ALL ^C	high	European	NA	D'Angelo et al, 2011 ³⁵
167 ALL ^C	high	European	NA	Erculj et al, 2012 ²²
186 ALL ^B	low	European	NA	Costea et al, 2006 ²⁸
201 LLA ^B	low	French-Canadian	NA	Krajnovic et al, 2004 ³³
240 LLA ^A	high	North American	NA	Kishi et al, 2007 ³⁴
520 LLA ^C	low	Various	NA	Aplenc et al, 2005 ²¹
64 ALL or ML ^A	high	European	-T	Faganel Kotnik et al, 2011 ²³
76 LLA ^C	high	Thai	-T	Pakakasama et al, 2007 ²⁰
81 LLA ^C	high	European	-T	Huang et al, 2008 ¹⁴
88 LLA ^C	high	European	-T	van Kooten et al, 2008 ³⁰
181 ALL ^B	high	Chinese	-T	Liu et al, 2011 ¹⁷
35 LLA ^C	high	Cretan	+T	Karathanasis et al, 2011 ¹⁹
37 LLA or NHL ^C	high	Turkish	+T	Kantar et al, 2009 ¹⁶
141 ALL ^C	high	Spanish	+T	Salazar et al, 2011 ²⁵

Type of sample: A: normal, B: tumor, C: unknown

High MTX dose = 1.5 – 5 g / m²; Low MTX dose = 15 – 30 mg / m²

NA, no association between the SNP and toxicity

+T, SNP is associated with increased toxicity (light shading)

-T, SNP is associated with decreased toxicity (dark shading)

Table 3. Types of toxicities analyzed and the findings in each study of the associations between the *MTHFR* C677T polymorphism and MTX toxicity.

MTHFR C677T									
Reference	Hematologic toxicity					MTX plasma levels	Mucositis	Hepatic toxicity	Other
	Anemia	Leucopenia	Neutropenia	Trombocytopenia	Myelosuppression				
Shimasaki et al, 2006 ¹⁸					NA		NA	NA	NA
Horinouchi et al, 2010 ²⁷								NA	
Karathanasis et al, 2011 ¹⁹	NA	NA		NA			NA	NA	
Chatzidakis et al, 2006 ²⁹	NA	NA						NA	
Kishi et al, 2003 ³⁶									NA
Pakakasama et al, 2007 ²⁰					NA		NA		NA
Huang et al, 2008 ¹⁴	NA	NA		NA		NA	NA	NA	NA
Lopez-Lopez et al, 2011 ¹⁵						NA			
Erculj et al, 2012 ²²	NA	NA		NA			NA	NA	NA
Kishi et al, 2007 ³⁴									NA
Krajinovic et al, 2004 ³³									NA
Aplenc et al, 2005 ²¹							NA	NA	NA
Kantar et al, 2009 ¹⁶	NA	NA		-T		NA		NA	NA
van Kooten et al, 2008 ³⁰		NA	-T	NA				NA	NA
Costea et al, 2006 ²⁸		-T	NA	NA				NA	
Shimasaki et al, 2008 ³⁷									+T
Imanishi et al, 2007 ⁴						+T		NA	
Tantawy et al, 2010 ²⁶			+T				+T	+T	+T
EL-Khodary et al, 2011 ³²	+T	+T		+T				+T	+T/NA
Faganel Kotnik et al, 2011 ²³		NA		NA			+T		NA
Salazar et al, 2011 ²⁵	NA	NA		+T			NA	NA	+T
D'Angelo et al, 2011 ³⁵					NA				+T
Liu et al, 2011 ¹⁷	NA		NA	+T	NA	NA	NA	NA	NA
Sepe et al, 2012 ³¹								+T	NA

NA, no association between the SNP and toxicity. +T, SNP is associated with increased toxicity. -T, SNP is associated with decreased toxicity

Table 4. Types of toxicities analyzed and the findings in each study of the associations between the *MTHFR* A1298C polymorphism and MTX toxicity.

MTHFR A1298C									
Reference	Hematologic toxicity					MTX plasma levels	Mucositis	Hepatic toxicity	Other
	Anemia	Leucopenia	Neutropenia	Trombocytopenia	Myelosuppression				
Tantawy et al, 2010 ²⁶			NA				NA	NA	NA
Lopez-Lopez et al, 2011 ¹⁵						NA			
D'Angelo et al, 2011 ³⁵					NA				NA
Erculj et al, 2012 ²²	NA	NA		NA			NA	NA	NA
Costea et al, 2006 ²⁸		NA	NA	NA				NA	
Krajinovic et al, 2004 ³³									NA
Kishi et al, 2007 ³⁴									NA
Aplenc et al, 2005 ²¹							NA	NA	NA
Faganel Kotnik et al, 2011 ²³		-T		NA			NA		NA
Pakakasama et al, 2007 ²⁰					-T		NA		NA
Huang et al, 2008 ¹⁴	NA	NA				NA	NA	NA	- T/NA
van Kooten et al, 2008 ³⁰		NA	NA	-T				NA	NA
Liu et al, 2011 ¹⁷	NA		NA	NA	NA	NA	NA	NA	-T
Karathanasis et al, 2011 ¹⁹	NA	NA		NA			NA	+T	
Kantar et al, 2009 ¹⁶	+T	NA		+T		+T		+T	+T
Salazar et al, 2011 ²⁵	NA	NA		+T			NA	NA	+T

NA, no association between the SNP and toxicity. +T, SNP is associated with increased toxicity. -T, SNP is associated with decreased toxicity