

AUTORREGRESIVE CONDITIONAL VOLATILITY, SKEWNESS AND KURTOSIS

Forthcoming in the Quarterly Journal of Economics and Finance

Ángel León
Universidad de Alicante
Dpto. Economía Financiera

Gonzalo Rubio
Universidad del País Vasco
Dpto. Fundamentos Análisis Económico II

Gregorio Serna
Universidad de Castilla – La Mancha
Dpto. Economía Financiera

December 2004

JEL Classification: G12, G13, C13, C14

Keywords: Conditional volatility, skewness and kurtosis; Gram-Charlier series expansion; Stock indices.

Corresponding author: Ángel León, Dpto. Economía Financiera, Facultad de Ciencia Económicas, Universidad de Alicante, Apartado de Correos 99, 03080 Alicante, Spain; E-mail: aleon@ua.es

We have received valuable comments from an anonymous referee, Alfonso Novales, Ignacio Peña and Hipolit Torró. Ángel León and Gonzalo Rubio acknowledge the financial support provided by the Ministerio de Ciencia y Tecnología, grants BEC2002-03797 and BEC2001-0636 respectively, and also thank the Fundación BBVA research grant 1-BBVA 00044.321-15466/2002

Abstract

This paper proposes a GARCH-type model allowing for time-varying volatility, skewness and kurtosis. The model is estimated assuming a Gram-Charlier series expansion of the normal density function for the error term, which is easier to estimate than the non-central t distribution proposed by Harvey and Siddique (1999). Moreover, this approach accounts for time-varying skewness and kurtosis while the approach by Harvey and Siddique (1999) only accounts for nonnormal skewness. We apply this method to daily returns of a variety of stock indices and exchange rates. Our results indicate a significant presence of conditional skewness and kurtosis. It is also found that specifications allowing for time-varying skewness and kurtosis outperform specifications with constant third and fourth moments.

AUTORREGRESIVE CONDITIONAL VOLATILITY, SKEWNESS AND KURTOSIS

1. Introduction

There have been many papers studying the departures from normality of asset return distributions. It is well known that stock return distributions exhibit negative skewness and excess kurtosis (see, for example, Harvey and Siddique, 1999; Peiró, 1999; and Premaratne and Bera, 2001). Specifically, excess kurtosis (the fourth moment of the distribution) makes extreme observations more likely than in the normal case, which means that the market gives higher probability to extreme observations than in normal distribution. However, the presence of negative skewness (the third moment of the distribution) has the effect of accentuating the left-hand side of the distribution. That is, the market gives higher probability to decreases than increases in asset pricing.

These issues have been widely analyzed in option pricing literature. For example, as explained by Das and Sundaram (1999), the well known volatility smile and smirk effects are closely related to the presence of excess kurtosis and negative skewness in the underlying asset returns distribution.

The generalized autorregressive conditional heteroscedasticity (GARCH) models, introduced by Engle (1982) and Bollerslev (1986), allow for time-varying volatility (but not for time-varying skewness or kurtosis). Harvey and Siddique (1999) present a way to jointly estimate time-varying conditional variance and skewness under a non-central t distribution for the error term in the mean equation. Their methodology is applied to several series of stock index returns, and it is found that autorregressive conditional skewness is significant and that the inclusion of skewness affects the persistence in variance. It is important to point out that the paper by Harvey and Siddique (1999) allows for time-varying skewness but still assumes constant kurtosis.

Premaratne and Bera (2001) have suggested capturing asymmetry and excess kurtosis with the Pearson type IV distribution, which has three parameters that can be interpreted as

volatility, skewness and kurtosis. This is an approximation to the non-central t distribution proposed by Pearson and Merrington (1958). However, these authors use time-varying conditional mean and variance, but maintain constant skewness and kurtosis over time. Similarly, Jondeau and Rockinger (2000) employ a conditional generalized Student-t distribution to capture conditional skewness and kurtosis by imposing a time-varying structure for the two parameters which control the probability mass in the assumed distribution¹. However, these parameters do not follow a GARCH structure for either skewness or kurtosis.

The purpose of this research is to extend the work by Harvey and Siddique (1999) assuming a distribution for the error term in the mean equation that accounts for nonnormal skewness and kurtosis. In particular, we jointly estimate time-varying volatility, skewness and kurtosis using a Gram-Charlier series expansion of the normal density function, along the lines suggested by Gallant and Tauchen (1989).

It is also worth noting that, apart from the fact that our approach accounts for time-varying kurtosis while the one by Harvey and Siddique (1999) does not, our likelihood function, based on a series expansion of the normal density function, is easier to estimate than the likelihood function based on the non-central t distribution employed by them.

The joint estimation of time-varying volatility, skewness and kurtosis can be useful in testing option pricing models that explicitly introduce the third and fourth moments of the underlying asset return distribution along the lines suggested by Heston (1993), Bates (1996), and Heston and Nandi (2000). It may also be useful in analyzing the information content of option-implied coefficients of skewness and kurtosis, extending the papers by Day and Lewis (1992), Lamoureux and Lastrapes (1993) and Amin and Ng (1997), among others.

The method proposed in this paper is applied to two different data sets. Firstly, our model is estimated using daily returns of four exchange rates series: British Pound/USD, Japanese

¹ This generalized Student-t distribution is based on Hansen's (1994) work.

Yen/USD, German Mark/USD and Swiss Franc/USD. Secondly we apply the method to five stock indices: S&P500 and NASDAQ100 (U.S.), DAX30 (Germany), IBEX35 (Spain), and the MEXBOL emerging market index (Mexico). These indices reflect the movements in their respective national financial markets and are used as underlying assets in several options and futures contracts.

Our results indicate significant presence of conditional skewness and kurtosis. It is also found that specifications allowing for time-varying skewness and kurtosis outperform specifications with constant third and fourth moments.

The rest of the paper is organized as follows. In Section 2 we present our GARCH-type model for estimating time-varying variance, skewness and kurtosis jointly. Section 3 presents the data and the empirical results regarding the estimation of the model. Section 4 compares the models allowing for time-varying skewness and kurtosis and the standard models with constant third and fourth moments. Section 5 concludes with a summary and discussion.

2. A model for conditional volatility, skewness and kurtosis

In this section we extend the model for conditional variance and skewness proposed by Harvey and Siddique (1999), to account for conditional kurtosis along the lines discussed in the introduction.

Given a series of asset prices $\{S_0, S_1, \dots, S_T\}$, we define continuously compounded returns for period t as $r_t = 100[\ln(S_t/S_{t-1})]$, $t = 1, 2, \dots, T$. Specifically, we present an asset return model containing either the GARCH(1,1) or NAGARCH (1,1) structure for conditional variance² and also a GARCH (1,1) structure for both conditional skewness and kurtosis. Under the NAGARCH specification for conditional variance, the model is denoted as

² Due to the well known leverage effect, we have chosen the NAGARCH (1,1) specification for the variance equation proposed by Engle and Ng (1993).

NAGARCHSK (and GARCHSK when conditional variance is driven by the GARCH (1,1) model³). It is given by:

$$\begin{aligned}
 r_t &= E_{t-1}(r_t) + \varepsilon_t; \quad \varepsilon_t \approx (0, \sigma_\varepsilon^2) \\
 \varepsilon_t &= h^{1/2} \eta_t; \quad \eta_t \approx (0, 1); \quad \varepsilon_t | I_{t-1} \approx (0, h_t) \\
 h_t &= \beta_0 + \beta_1 (\varepsilon_{t-1} + \beta_3 h_{t-1}^{1/2})^2 + \beta_2 h_{t-1} \\
 s_t &= \gamma_0 + \gamma_1 \eta_{t-1}^3 + \gamma_2 s_{t-1} \\
 k_t &= \delta_0 + \delta_1 \eta_{t-1}^4 + \delta_2 k_{t-1}
 \end{aligned} \tag{1}$$

where $E_{t-1}(\bullet)$ denotes the conditional expectation on an information set till period $t-1$ denoted as I_{t-1} . We establish that $E_{t-1}(\eta_t) = 0$, $E_{t-1}(\eta_t^2) = 1$, $E_{t-1}(\eta_t^3) = s_t$ and $E_{t-1}(\eta_t^4) = k_t$ where both s_t and k_t are driven by a GARCH (1,1) structure. Hence, s_t and k_t represent respectively skewness and kurtosis corresponding to the conditional distribution of the standardized residual $\eta_t = \varepsilon_t h_t^{-1/2}$.

Using a Gram-Charlier (GC) series expansion of the normal density function and truncating at the fourth moment⁴, we obtain the following density function for the standardized residuals η_t conditional on the information available in $t-1$:

$$\begin{aligned}
 g(\eta_t | I_{t-1}) &= \phi(\eta_t) \left[1 + \frac{s_t}{3!} (\eta_t^3 - 3\eta_t) + \frac{k_t - 3}{4!} (\eta_t^4 - 6\eta_t^2 + 3) \right] \\
 &= \phi(\eta_t) \psi(\eta_t)
 \end{aligned} \tag{2}$$

³ Specifically, in the equations below, we obtain the GARCHSK model for $\beta_3 = 0$.

where $\phi(\bullet)$ denotes the probability density function (henceforth pdf) corresponding to the standard normal distribution and $\Psi(\bullet)$ is the polynomial part of fourth order corresponding to the expression between brackets in (2). Note that the pdf defined in (2) is not really a density function because for some parameter values in (1) the density $g(\bullet)$ might be negative due to the component $\Psi(\bullet)$. Similarly, the integral of $g(\bullet)$ on \mathfrak{R} is not equal to one. We propose a true pdf, denoted as $f(\bullet)$, by transforming the density $g(\bullet)$ according to the method in Gallant and Tauchen (1989). Specifically, in order to obtain a well defined density everywhere we square the polynomial part $\Psi(\bullet)$, and to insure that the density integrates to one we divide by the integral of $g(\bullet)$ over \mathfrak{R} ⁵. The resulting pdf written in abbreviated form is⁶:

$$f(\eta_t | I_{t-1}) = \phi(\eta_t) \psi^2(\eta_t) / \Gamma_t \quad (3)$$

where

$$\Gamma_t = 1 + \frac{s_t^2}{3!} + \frac{(k_t - 3)^2}{4!}$$

Therefore, after omitting unessential constants, the logarithm of the likelihood function for one observation corresponding to the conditional distribution $\varepsilon_t = h_t^{1/2} \eta_t$, whose pdf is $h_t^{-1/2} f(\eta_t | I_{t-1})$, is given by

$$l_t = -\frac{1}{2} \ln h_t - \frac{1}{2} \eta_t^2 + \ln(\psi^2(\eta_t)) - \ln(\Gamma_t) \quad (4)$$

As pointed out before, this likelihood function is clearly easier to estimate than the one based on a non-central t proposed by Harvey and Siddique (1999). In fact, the likelihood function in (4) is the same as in the standard normal case plus two adjustment terms accounting for time-varying skewness and kurtosis. Moreover, it is worth noting that the

⁴ See Jarrow and Rudd (1982) and also Corrado and Su (1996).

⁵ See the appendix for proof that this nonnegative function is really a density function that integrates to one.

⁶ An alternative approach under the Gram-Charlier framework is proposed by Jondeau and Rockinger (2001) who also show how constraints on the parameters defining skewness and kurtosis may be implemented to insure that the expansion defines a density. However, their approach does not seem to be feasible in both skewness and kurtosis within the conditional case.

density function based on a Gram-Charlier series expansion in equation (3) nests the normal density function (when $s_t = 0$ and $k_t = 3$), while the noncentral t does not. Therefore, the restrictions imposed by the normal density function with respect to the more general density based on a Gram-Charlier series expansion can be easily tested. Finally, note that NAGARCHSK nests the GARCH (1,1) specification for the conditional variance when $\beta_3 = 0$ in (1). We denote this nested case as the GARCHSK model.

3. Empirical results

3.1 Data and preliminary findings

Our methodology is applied to two different data sets. The first one includes daily returns of five exchange rates series: British Pound/USD (GBP/USD), Japanese Yen/USD (JPY/USD), German Mark/USD (GEM/USD) and Swiss Franc/USD (CHF/USD). The second data set includes five stock indexes: S&P500 and NASDAQ100 (U.S.), DAX30 (Germany), IBEX35 (Spain) and the emerging market index MEXBOL (Mexico).

Our data set includes daily closing prices from January 2, 1990 to May 3, 2002 for the five exchange rate series, and from January 2, 1990 to July 17, 2003 for all stock index series except for MEXBOL, which includes data from January 2, 1995 to July 17, 2003. These closing prices are employed to calculate the corresponding continuously compounded daily returns, and Table 1 presents some descriptive statistics. Note that all series show leptokurtosis and there is also evidence of negative skewness except for GBP/USD and MEXBOL. It is also worth noting that the Mexican emerging market returns (MEXBOL) show the highest values of unconditional standard deviation, skewness and kurtosis.

Before we estimate our NAGARCHSK model, we analyze the dynamic structure in the mean equation of (1). Specifically, the ARMA structure that maximizes the Schwarz Information Criterion (SIC) is selected. All the parameters implied in every model below are estimated by maximum likelihood assuming that the Gram-Charlier series expansion distribution given by (3) holds for the error term, and using Bollerslev and Wooldridge

(1992) robust standard errors⁷. If we define the SIC as $\ln(L_{ML}) - (q/2)\ln(T)$, where q is the number of estimated parameters, T is the number of observations, and L_{ML} is the value of the log likelihood function using the q estimated parameters, then the selected model is the one with the highest SIC. According to SIC, MA(1) and AR(1) models without constant term yield very similar results⁸. However, the AR(1) has the advantage of being consistent with the nonsynchronous contracts of individual stocks which constitute the indices. Definitively, the dynamic conditional mean structure for every estimation is represented by an AR(1) model with no constant term.

Table 2 presents the Ljung-Box statistics of order 20, denoted as LB(20), for ε_t^2 , ε_t^3 and ε_t^4 , where ε_t is the error term in the AR(1) model (with no constant term). The statistic for all moments is quite large (p-value = 0.000 in all cases). In other words, the significant serial correlation for ε_t^2 , ε_t^3 and ε_t^4 indicates time-varying volatility, skewness and kurtosis, and it justifies the estimation of our GARCHSK or NAGARCHSK models defined in (1) with time-varying volatility, skewness and kurtosis.

3.2 Model estimation with time-varying volatility, skewness and kurtosis

Before presenting the estimation results obtained with both the exchange rates and the stock indexes series, we summarize the four nested models estimated as follows:

$$\text{Mean:} \quad r_t = \alpha_1 r_{t-1} + \varepsilon_t \quad (5-a)$$

$$\text{Variance (GARCH):} \quad h_t = \beta_0 + \beta_1 \varepsilon_{t-1}^2 + \beta_2 h_{t-1} \quad (5-b)$$

$$\text{Variance (NAGARCH):} \quad h_t = \beta_0 + \beta_1 \left(\varepsilon_{t-1} + \beta_3 h_{t-1}^{1/2} \right)^2 + \beta_2 h_{t-1} \quad (5-c)$$

$$\text{Skewness:} \quad s_t = \gamma_0 + \gamma_1 \varepsilon_{t-1}^3 + \gamma_2 s_{t-1} \quad (5-d)$$

$$\text{Kurtosis:} \quad k_t = \delta_0 + \delta_1 \varepsilon_{t-1}^4 + \delta_2 k_{t-1} \quad (5-e)$$

⁷ All maximum likelihood estimations in this paper are carried out using the CML subroutine of GAUSS.

⁸ The constant terms were never significant in previous tests.

We estimate first two standard models for conditional variance: the GARCH (1,1) model (equations (5-a) and (5-b)), and the NAGARCH (1,1) model (equations (5-a) and (5-c)), where a normal distribution is assumed for the unconditional standardized error η_t . We also estimate the generalizations of the standard GARCH and NAGARCH models, with time-varying skewness and kurtosis, named GARCHSK (equations (5-a), (5-b), (5-d) and (5-e)) and NAGARCHSK (equations (5-a), (5-c), (5-d) and (5-e)), assuming in both cases the distribution based on the Gram-Charlier series expansion given by equation (3). In the NAGARCH specification of the variance equation, a negative value of β_3 implies a negative correlation between shocks and conditional variance.

It should be noted that, given that the likelihood function is highly nonlinear, special care must be taken in selecting the starting values of the parameters. As usual in these cases, given that the four models are nested, the estimation is performed following several stages, and using the parameters estimated from the simpler models as starting values for more complex ones.

The results for the exchange rate series are presented in Tables 3 and 4 for the GARCH and GARCHSK models respectively. It is found that for all exchange rates series the coefficient for asymmetric variance, β_3 , is not significant, confirming that the leverage effect, commonly observed in other financial series, is not observed in the case of exchange rates. Therefore, for the exchange rate series only the results for symmetric variance models are presented.

As expected, the results for all exchange rate series indicate a significant presence of conditional variance. Volatility is found to be persistent since the coefficient of lagged volatility is positive and significant, indicating that high conditional variance is followed by high conditional variance.

Moreover, it is found that for the GBP/USD, DEM/USD and CHF/USD exchange rate series, days with high skewness are followed by days with high skewness, since the coefficient for lagged skewness (γ_2) is positive and significant, although its magnitude is

lower than in the variance case. Also, shocks to skewness are significant, although they are less relevant than its persistence. However, there seems to be no structure in skewness in the JPY/USD series, since neither γ_1 nor γ_2 is significant in this case.

As with skewness, the results for the kurtosis equation indicate that days with high kurtosis are followed by days with high kurtosis, since the coefficient for lagged kurtosis (δ_2) is positive and significant. Its magnitude is greater than that of skewness but still lower than that of variance. As before, shocks to kurtosis are significant, except for the JPY/USD series.

Finally, it is worth noting that the value of the SIC, shown at the bottom of Tables 3 and 4, rises monotonically in all cases when we move from the simpler models to the more complicated ones, with the GARCHSK model showing the highest figure. Therefore, for the four exchange rates series analyzed, the GARCHSK specification seems to be the most appropriate one according to the SIC criterion.

The results for the five stock indices are presented in Tables 5, 6, 7 and 8 for GARCH, NAGARCH, GARCHSK and NAGARCHSK models respectively.

As expected, the results shown in Table 5 (GARCH models) indicate significant presence of conditional variance, with the two American indices (S&P500 and NASDAQ100) showing the highest degree of persistence. However, Table 6 (NAGARCH models) shows that contrary to the exchange rate case, the coefficient for asymmetric variance, β_3 , is negative and significant, confirming the presence of the leverage effect commonly observed in the markets.

In regard to the skewness equation (Tables 7 and 8), as before, significant presence of conditional skewness is found, with at least one of the coefficients associated with shocks to skewness (γ_1) and to lagged skewness (γ_2) being significant, except for S&P500 stock index under the NAGARCHSK specification.

Similar results are obtained for the kurtosis equation with both GARCHSK and NAGARCSK specifications. The coefficient associated with shocks to kurtosis (δ_1) is significant in all cases, except for NASDAQ100 with the GARCHSK model and to some extent for IBEX35 with the NAGARCH model. Moreover, the coefficient associated with lagged kurtosis (δ_2) is significant in all cases except S&P500 with both specifications. Nevertheless, there is significant presence of conditional kurtosis for all stock indices, with both specifications, since at least one of the coefficients associated with shocks to kurtosis or to lagged kurtosis is found to be significant.

As obtained with the exchange rate series, the value of the SIC rises monotonically for all stock index series analyzed when we move from the simpler models to the more complicated ones, with the NAGARCHSK model showing the highest value. This seems to be the most appropriate specification.

4. Comparing the models

One way to start comparing the models is to compute a likelihood ratio test. It is easy to see that the density function based on a Gram-Charlier series expansion in equation (3) nests the normal density function when $s_t = 0$ and $k_t = 3$ (alternatively when $\gamma_1 = \gamma_2 = \gamma_3 = 0$, $\delta_1 = 3$ and $\delta_2 = \delta_3 = 0$). Therefore, the restrictions imposed by the normal density function with respect to the more general density based on a Gram-Charlier series expansion can be tested by means of a likelihood ratio test. The results are contained in Table 9. The value of the LR statistic is quite large in all cases, indicating the rejection of the null hypothesis that the true density is the restricted one, i.e. the normal density function.

A second way of comparing the models is to compare the properties of the conditional variances obtained with each model. Figure 1 shows the behavior of conditional variance for one of the exchange rate series -GBP/USD- with both GARCH and GARCHSK models, and for one of the stock index series -S&P500- with both NAGARCH and NAGARCHSK specifications. It is clear that conditional variances obtained with models accounting for time-varying skewness and kurtosis are smoother than those obtained with standard

GARCH or NAGARCH models. This is confirmed by the results in Table 10, which shows some descriptive statistics for these conditional variances. In fact, conditional variances obtained with GARCHSK or NAGARCHSK models show less standard deviation, skewness and kurtosis than those obtained with the standard models. This fact was observed by Harvey and Siddique (1999) with their time-varying skewness (although constant-kurtosis) specification.

The in-sample predictive ability of the different models is compared by means of two metrics. The variable predicted is the squared forecast error (ε_t^2) and the predictors are the conditional variances (h_t) from, respectively, the standard GARCH or NAGARCH models and GARCHSK or NAGARCHSK models. The two metrics are:

Median absolute error: $MAE = med(|\varepsilon_t^2 - h_t|)$

Median percentage absolute error: $MPAE = med\left(\frac{|\varepsilon_t^2 - h_t|}{\varepsilon_t^2}\right)$

The metrics are based on the median since it is more robust than the mean in view of the high dispersal of the error series. The results are shown in Table 11. Models accounting for time-varying skewness and kurtosis outperform standard GARCH or NAGARCH models. They are the best performing models with the two metrics with all exchange rates and stock index series except for NASDAQ100 and IBEX35 with the median absolute error (although not with the median percentage absolute error).

Furthermore, it is worth noting that the series that performs best, based on these metrics, is the MEXBOL stock index, which is the series with the highest values of unconditional standard deviation, skewness and kurtosis (Table 1). This result could suggest the potential application of our methodology to financial series from emerging economies, characterized by higher risk and more pronounced departures from normality.

5. Conclusions

It is well known that the generalized autorregressive conditional heteroscedasticity (GARCH) models, introduced by Engle (1982) and Bollerslev (1986) allow for time-varying volatility (but not for time-varying skewness or kurtosis). However, given the increasing attention that time-varying skewness and kurtosis have attracted in option pricing literature, it may be useful to analyze a model that jointly accounts for conditional second, third and fourth moments.

Harvey and Siddique (1999) present a way of jointly estimating time-varying conditional variance and skewness, assuming a non-central t distribution for the error term in the mean equation. We propose a GARCH-type model allowing for time-varying volatility, skewness and kurtosis. The model is estimated assuming a Gram-Charlier series expansion of the normal density function, along the lines suggested by Gallant and Tauchen (1989), for the error term in the mean equation. This distribution is easier to estimate than the non-central t distribution proposed by Harvey and Siddique (1999). Moreover, our approach accounts for time-varying skewness and kurtosis while the one by Harvey and Siddique (1999) only accounts for time-varying skewness.

Firstly, our model is estimated using daily returns of four exchange rate series, five stock indices and the emerging market index MEXBOL (Mexico). Our results indicate significant presence of conditional skewness and kurtosis. Moreover, it is found that specifications allowing for time-varying skewness and kurtosis outperform specifications with constant third and fourth moments.

Finally, it is important to point out two main implications of our GARCHSK and NAGARCHSK model. First, they can be useful in estimating future coefficients of volatility, skewness and kurtosis, which are unknown parameters in option pricing models that account for nonnormal skewness and kurtosis. For example, estimates of volatility, skewness and kurtosis from the NAGARCHSK model, based on historical series of returns, could be compared with option implied coefficients in terms of their out of sample option pricing performance. Secondly, our models could be useful in testing the information

content of option implied coefficients of volatility, skewness and kurtosis. This could be done by including option implied coefficients as exogenous terms in the equations of volatility, skewness and kurtosis, extending the papers by Day and Lewis (1992), Lamoureux and Lastrapes (1993) and Amin and Ng (1997), among others.

APPENDIX

Here we show that the nonnegative function $f(\eta_t|I_{t-1})$ in (3) is really a density function, that is it integrates to one. We can rewrite $\psi(\eta_t)$ in (2) as:

$$\psi(\eta_t) = 1 + \frac{s_t}{\sqrt{3!}} H_3(\eta_t) + \frac{k_t - 3}{\sqrt{4!}} H_4(\eta_t)$$

where $\{H_i(x)\}_{i \in \mathbb{N}}$ represents the Hermite polynomials such that $H_0(x) = 1$, $H_1(x) = x$ and for $i \geq 2$ they hold the following recurrence relation:

$$H_i(x) = (xH_{i-1}(x) - \sqrt{i-1}H_{i-2}(x)) / \sqrt{i}$$

It is verified that $\{H_i(x)\}_{i \in \mathbb{N}}$ is an orthonormal basis satisfying that:

$$\int_{-\infty}^{\infty} H_i(x)\phi(x)dx = 1, \quad \forall i \tag{A-1}$$

$$\int_{-\infty}^{\infty} H_i(x)H_j(x)\phi(x)dx = 0, \quad \forall i \neq j \tag{A-2}$$

where $\phi(\bullet)$ denotes the $N(0,1)$ density function. If we integrate the conditional density function in (3), given conditions (A-1) and (A-2):

$$\begin{aligned} & (1/\Gamma_t) \int_{-\infty}^{\infty} \phi(\eta_t) \left[1 + \frac{s_t}{\sqrt{3!}} H_3(\eta_t) + \frac{k_t - 3}{\sqrt{4!}} H_4(\eta_t) \right]^2 d\eta_t \\ &= (1/\Gamma_t) \left[\int_{-\infty}^{\infty} \phi(\eta_t) d\eta_t + \frac{s_t^2}{3!} \int_{-\infty}^{\infty} H_3^2(\eta_t)\phi(\eta_t) d\eta_t + \frac{(k_t - 3)^2}{4!} \int_{-\infty}^{\infty} H_4^2(\eta_t)\phi(\eta_t) d\eta_t \right] \\ &= (1/\Gamma_t) \left[1 + \frac{s_t^2}{3!} + \frac{(k_t - 3)^2}{4!} \right] \\ &= 1. \end{aligned}$$

REFERENCES

- Amin, K. and V. Ng (1997), "Inferring Future Volatility from the Information in Implied Volatility in Eurodollar Options: A New Approach", *The Review of Financial Studies* 10, 333-367.
- Backus, D., Foresi, S., Li, K. and L. Wu (1997), "Accounting for Biases in Black-Scholes", *Working Paper, Stern School of Business, New York University*.
- Bates, D. (1996), "Jumps and Stochastic Volatility: Exchange Rate Processes Implicit in Deutschemark Options", *The Review of Financial Studies* 9, 69-107.
- Bollerslev, T. (1986), "Generalized Autorregresive Conditional Heteroskedasticity", *Journal of Econometrics* 31, 307-327.
- Bollerslev, T. and J. Wooldridge (1992), "Quasi-Maximum Likelihood Estimation and Inference y Dynamic Models with Time-Varying Covariances", *Econometric Reviews* 11, 143-172.
- Corrado, C. and T. Su (1996), "Skewness and Kurtosis in S&P 500 index returns implied by option prices", *Journal of Financial Research* 19, 175-192.
- Das, S and R. Sundaram (1999), "Of Smiles and Smirks: A Term Structure Perspective", *Journal of Financial and Quantitative Analysis* 34, 211-239.
- Day, T. and C. Lewis (1992), "Stock market volatility and the information content of stock index options", *Journal of Econometrics* 52, 267-287.
- Engle, R. F. (1982), "Autorregresive Conditional Heteroskedasticity with Estimates of the Variance of U.K. Inflation", *Econometrica* 50, 987-1008.

Engle, R. F. and V. K. Ng (1993), "Measuring and Testing the Impact of News on Volatility", *The Journal of Finance* 48, 1749-1778.

Gallant, A. R. and G. Tauchen (1989), "Seminonparametric Estimation of Conditionally Constrained Heterogeneous Processes: Asset Pricing Applications", *Econometrica* 57, 1091-1120.

Hansen, B. (1994), "Autoregressive Conditional Density Estimation", *International Economic Review* 35, 705-730.

Harvey, C. R. and A. Siddique (1999), "Autoregressive Conditional Skewness", *Journal of Financial and Quantitative Analysis* 34, 465-487.

Heston, S. (1993), "A Closed-Form Solution for Options with Stochastic Volatility, with Applications to Bond and Currency Options", *The Review of Financial Studies* 6, 327-343.

Heston, S. and S. Nandi (2000), "A Closed-Form GARCH Option Valuation Model", *The Review of Financial Studies* 13, 585-625.

Jarrow, R. and A. Rudd (1982), "Approximate Option Valuation for Arbitrary Stochastic Processes", *Journal of Financial Economics* 10, 347-369.

Jondeau, E. and M. Rockinger (2000), "Conditional Volatility, Skewness and Kurtosis: Existence and Persistence", *Working Paper, HEC School of Management*.

Jondeau, E. and M. Rockinger (2001), "Gram-Charlier Densities", *Journal of Economic Dynamics & Control* 25, 1457-1483.

Lamoureux, C. and W. Lastrapes (1993), "Forecasting Stock-Return Variance: Toward an Understanding of Stochastic Implied Volatilities", *The Review of Financial Studies* 6, 293-326.


Pearson, E. S. and M. Merrington (1958), "An approximation to the distribution of noncentral t ", *Biometrika* 45, 484-491.

Peiró, A. (1999), "Skewness in financial returns", *Journal of Banking and Finance* 23, 847-862.


Premaratne, G. and A. K. Bera (2001), "Modeling Asymmetry and Excess Kurtosis in Stock Return Data", *Working Paper, Department of Economics, University of Illinois*.

FIGURE 1
ESTIMATED CONDITIONAL VARIANCES WITH NAGARCH AND NAGARCHSK
MODELS


CONDITONAL VARIANCE
GARCH GBP/USD



CONDITONAL VARIANCE
GARCHSK GBP/USD



CONDITONAL VARIANCE
NAGARCH S&P500



CONDITONAL VARIANCE
NAGARCHSK S&P500




TABLE 1
DESCRIPTIVE STATISTICS FOR DAILY RETURNS

PANEL A: EXCHANGE RATES				
STATISTIC	GBP/USD	JPY/USD	DEM/USD	CHF/USD
Simple size	3126	3126	3126	3126
Mean	0.0030	-0.0045	0.0072	0.0003
Median	0.0000	0.0120	0.0207	0.0217
Maximum	3.2860	3.3004	3.1203	3.0747
Minimum	-2.8506	-5.7093	-2.9497	-3.7243
Stand. Dev.	0.5731	0.7192	0.6621	0.7197
Skewness	0.2334	-0.5794	-0.0594	-0.2000
Kurtosis	5.7502	7.3298	4.6546	4.5432
Jarque-Bera (p-value)	1013.565 (0.0000)	2616.775 (0.0000)	358.4119 (0.0000)	331.0593 (0.0000)

PANEL B: STOCK INDEXES					
STATISTIC	S&P500	NASDAQ	DAX30	IBEX35	MEXBOL
Simple size	3415	3416	3407	3390	2137
Mean	0.0294	0.0383	0.0178	0.0246	0.0511
Median	0.0315	0.1217	0.0641	0.0508	0.0099
Maximum	5.5732	13.2546	7.5527	6.8372	12.1536
Minimum	-7.1127	-10.1684	-8.8747	-8.8758	-14.3139
Stand. Dev.	1.0611	1.6117	1.5056	1.3876	1.8086
Skewness	-0.0995	-0.0099	-0.1944	-0.1854	0.0712
Kurtosis	6.5658	8.3740	6.3210	5.9169	8.6060
Jarque-Bera (p-value)	1814.880 (0.0000)	4110.566 (0.0000)	1587.134 (0.0000)	1221.204 (0.0000)	2800.124 (0.0000)

TABLE 2
LJUNG-BOX STATISTICS WITH ORDER 20 OF RESIDUALS FROM AR(1) MODEL

The table presents the Ljung-Box statistic (asymptotic p-value in parenthesis) with order 20, i.e. LB(20), of ε_t^2 , ε_t^3 and ε_t^4 , where ε_t is the error term from an AR(1) model for daily returns (in bold are significantly different from zero Ljung-Box statistics)

SERIES	LB(20) - ε_t^2	LB(20) - ε_t^3	LB(20) - ε_t^4
GBP/USD	825.43 (0.000)	134.37 (0.000)	332.34 (0.000)
JPY/USD	567.01 (0.000)	208.55 (0.000)	196.37 (0.000)
DEM/USD	407.25 (0.000)	70.501 (0.000)	187.38 (0.000)
CHF/USD	317.69 (0.000)	133.75 (0.000)	365.89 (0.000)
S&P500	131.81 (0.000)	120.91 (0.000)	139.79 (0.000)
NASDAQ	3152.1 (0.000)	252.04 (0.000)	315.26 (0.000)
DAX30	2919.1 (0.000)	72.889 (0.000)	489.37 (0.000)
IBEX35	1719.1 (0.000)	131.16 (0.000)	271.49 (0.000)
MEXBOL	488.67 (0.000)	238.18 (0.000)	283.82 (0.000)

TABLE 3
GARCH MODELS – EXCHANGE RATES

The reported coefficients shown in each row of the table are ML estimates of the standard GARCH model:

$$r_t = \alpha_1 r_{t-1} + \varepsilon_t$$

$$h_t = \beta_0 + \beta_1 \varepsilon_{t-1}^2 + \beta_2 h_{t-1}$$

for percentage daily returns of British Pound/American Dollar (GBP/USD), Japanese Yen/US Dollar (JPY/USD), German Mark/US Dollar (DEM/USD) and Swiss Franc/US Dollar (CHF/USD) exchange rates, from January 1990 to March 2002. $h_t = \text{var}(r_t | r_{t-1}, r_{t-2}, \dots)$, $\varepsilon_t | \varepsilon_{t-1}, \varepsilon_{t-2}, \dots$ follows a $N(0, h_t)$ distribution. All models have been estimated by ML using the Berndt-Hall-Hall-Hausman algorithm (quasi-maximum likelihood p-values in parenthesis; in bold are significantly different from zero coefficients at 5%).

	Parameter	GBP/USD	JPY/USD	DEM/USD	CHF/USD
Mean equation	α_1	0.0432 (0.0263)	0.0175 (0.3826)	0.0364 (0.0573)	0.0304 (0.1154)
Variance equation	β_0	0.0031 (0.0459)	0.0086 (0.0645)	0.0051 (0.0663)	0.0111 (0.0715)
	β_1	0.0435 (0.0000)	0.0428 (0.0011)	0.0378 (0.0000)	0.0336 (0.0003)
	β_2	0.9468 (0.0000)	0.9402 (0.0000)	0.9502 (0.0000)	0.94445 (0.0000)
Log-Likelihood	-	409.3328	-352.5956	-149.3089	-451.7276
SIC	-	393.2391	-368.6843	-165.4027	-467.8213

TABLE 4
GARCHSK MODELS – EXCHANGE RATES

The reported coefficients shown in each row of the table are ML estimates of the GARCHSK model:

$$r_t = \alpha_1 r_{t-1} + \varepsilon_t$$

$$h_t = \beta_0 + \beta_1 \varepsilon_{t-1}^2 + \beta_2 h_{t-1}$$

$$s_t = \gamma_0 + \gamma_1 \eta_{t-1}^3 + \gamma_2 s_{t-1}$$

$$k_t = \delta_0 + \delta_1 \eta_{t-1}^4 + \delta_2 k_{t-1}$$

for percentage daily returns of of British Pound/US Dollar (GBP/USD), Japanese Yen/US Dollar (JPY/USD), German Mark/US Dollar (DEM/USD) and Swiss Franc/US Dollar (CHF/USD) exchange rates, from January 1990 to March 2002. $h_t = \text{var}(r_t | r_{t-1}, r_{t-2}, \dots)$, $s_t = \text{skewness}(r_t | r_{t-1}, r_{t-2}, \dots)$, $k_t = \text{kurtosis}(r_t | r_{t-1}, r_{t-2}, \dots)$, $\eta_t = \varepsilon_t h_t^{-1/2}$, and $\varepsilon_t | \varepsilon_{t-1}, \varepsilon_{t-2}, \dots$ follows the distribution based on a Gram-Charlier series expansion. All models have been estimated by ML using the Berndt-Hall-Hall-Hausman algorithm (quasi-maximum likelihood p-values in parenthesis; in bold are significantly different from zero coefficients at 5%).

	Parameter	GBP/USD	JPY/USD	DEM/USD	CHF/USD
Mean equation	α_1	0.0219 (0.2537)	-0.0030 (0.8670)	0.0249 (0.3804)	0.0015 (0.9322)
	β_0	0.0015 (0.0783)	0.0061 (0.0378)	0.0022 (0.0159)	0.0075 (0.0007)
Variance equation	β_1	0.0366 (0.0000)	0.0309 (0.0021)	0.0236 (0.0000)	0.0217 (0.0000)
	β_2	0.9550 (0.0000)	0.9537 (0.0000)	0.9690 (0.0000)	0.9611 (0.0000)
	γ_0	0.0053 (0.5379)	-0.0494 (0.0482)	-0.0270 (0.0398)	-0.0242 (0.0989)
Skewness equation	γ_1	0.0093 (0.0004)	0.0018 (0.4190)	0.0175 (0.0054)	0.0054 (0.0688)
	γ_2	0.6180 (0.0000)	0.3414 (0.2097)	0.4421 (0.0000)	0.6468 (0.0002)
	δ_0	1.3023 (0.0000)	1.2365 (0.0038)	1.9649 (0.0000)	0.5500 (0.0000)
Kurtosis equation	δ_1	0.0028 (0.0000)	0.0014 (0.1102)	0.01356 (0.0000)	0.0060 (0.0000)
	δ_2	0.6229 (0.0000)	0.6464 (0.0000)	0.4045 (0.0002)	0.8303 (0.0000)
	Log-Likelihood	-	472.3652	-237.6668	-117.5896
SIC	-	432.1309	-277.9012	-157.8240	-461.2317

TABLE 5
GARCH MODELS - STOCK INDICES

The reported coefficients shown in each row of the table are ML estimates of the standard GARCH model:

$$r_t = \alpha_1 r_{t-1} + \varepsilon_t$$

$$h_t = \beta_0 + \beta_1 \varepsilon_{t-1}^2 + \beta_2 h_{t-1}$$

for percentage daily returns of S&P500, NASDAQ100, DAX30, IBEX35 stock indices, from January 1990 to July 2003, and MEXBOL from January 1995 to July 2003. $h_t = \text{var}(r_t | r_{t-1}, r_{t-2}, \dots)$, $\varepsilon_t | \varepsilon_{t-1}, \varepsilon_{t-2}, \dots$ follows a $N(0, h_t)$ distribution. All models have been estimated by ML using the Berndt-Hall-Hall-Hausman algorithm (quasi-maximum likelihood p-values in parenthesis; in bold are significantly different from zero coefficients at 5%).

	Parameter	S&P500	NASDAQ	DAX30	IBEX35	MEXBOL
Mean equation	α_1	0.03394 (0.0544)	0.1266 (0.0000)	0.0179 (0.3133)	0.0943 (0.0000)	0.1564 (0.0000)
Variance equation	β_0	0.0055 (0.0414)	0.0149 (0.0155)	0.0317 (0.0092)	0.05741 (0.0026)	0.0827 (0.0958)
	β_1	0.0587 (0.0000)	0.0948 (0.0000)	0.09394 (0.0000)	0.1035 (0.0000)	0.1194 (0.0098)
	β_2	0.9379 (0.0000)	0.9009 (0.0000)	0.8918 (0.0000)	0.8666 (0.0000)	0.8591 (0.0000)
Log-Likelihood	-	-1459.6826	-2424.1550	-2525.9824	-2441.0090	-2095.6885
SIC	-	-1475.9532	-2440.4262	-2542.2484	-2457.2650	-2111.0210

TABLE 6
NAGARCH MODELS – STOCK INDICES

The reported coefficients shown in each row of the table are ML estimates of the NAGARCH model:

$$r_t = \alpha_1 r_{t-1} + \varepsilon_t$$

$$h_t = \beta_0 + \beta_1 (\varepsilon_{t-1} + \beta_3 h_{t-1}^{1/2})^2 + \beta_2 h_{t-1}$$

for percentage daily returns of S&P500, NASDAQ100, DAX30, IBEX35 stock indices, from January 1990 to July 2003, and MEXBOL from January 1995 to July 2003. $h_t = \text{var}(r_t | r_{t-1}, r_{t-2}, \dots)$, $\varepsilon_t | \varepsilon_{t-1}, \varepsilon_{t-2}, \dots$ follows a $N(0, h_t)$ distribution. All models have been estimated by ML using the Berndt-Hall-Hausman algorithm (quasi-maximum likelihood p-values in parenthesis; in bold are significantly different from zero coefficients at 5%).

	Parameter	S&P500	NASDAQ	DAX30	IBEX35	MEXBOL
Mean equation	α_1	0.0461 (0.0098)	0.1387 (0.0098)	0.0200 (0.2602)	0.0956 (0.0000)	0.1665 (0.0000)
Variance equation	β_0	0.0126 (0.0028)	0.0270 (0.0055)	0.0332 (0.0010)	0.0560 (0.0009)	0.0852 (0.0142)
	β_1	0.0607 (0.0000)	0.1086 (0.0000)	0.0758 (0.0000)	0.0865 (0.0000)	0.0961 (0.0004)
	β_2	0.8776 (0.0000)	0.8605 (0.0000)	0.8855 (0.0000)	0.8609 (0.0000)	0.8169 (0.0000)
	β_3	-0.9588 (0.0000)	-0.4828 (0.0000)	-0.5678 (0.0000)	-0.5326 (0.0000)	-0.8349 (0.0000)
Log-Likelihood	-	-1401.8598	-2385.3512	-2496.0414	-2413.6763	-2050.0510
SIC	-	-1422.1982	-2405.6903	-2516.3739	-2433.9963	-2069.2165

TABLE 7
GARCHSK MODELS – STOCK INDICES

The reported coefficients shown in each row of the table are ML estimates of the GARCHSK model:

$$r_t = \alpha_1 r_{t-1} + \varepsilon_t$$

$$h_t = \beta_0 + \beta_1 \varepsilon_{t-1}^2 + \beta_2 h_{t-1}$$

$$s_t = \gamma_0 + \gamma_1 \eta_{t-1}^3 + \gamma_2 s_{t-1}$$

$$k_t = \delta_0 + \delta_1 \eta_{t-1}^4 + \delta_2 k_{t-1}$$

for percentage daily returns of S&P500, NASDAQ100, DAX30, IBEX35 stock indices, from January 1990 to July 2003, and MEXBOL from January 1995 to July 2003. $h_t = \text{var}(r_t | r_{t-1}, r_{t-2}, \dots)$, $s_t = \text{skewness}(r_t | r_{t-1}, r_{t-2}, \dots)$, $k_t = \text{kurtosis}(r_t | r_{t-1}, r_{t-2}, \dots)$, $\eta_t = \varepsilon_t h_t^{-1/2}$, and $\varepsilon_t | \varepsilon_{t-1}, \varepsilon_{t-2}, \dots$ follows the distribution based on a Gram-Charlier series expansion. All models have been estimated by ML using the Berndt-Hall-Hall-Hausman algorithm (quasi-maximum likelihood p-values in parenthesis; in bold are significantly different from zero coefficients at 5%).

	Parameter	S&P500	NASDAQ	DAX30	IBEX35	MEXBOL
Mean equation	α_1	0.0211	0.1229	0.0080	0.0949	0.1775
		(0.2285)	(0.0000)	(0.6557)	(0.0000)	(0.0000)
Variance equation	β_0	0.0023	0.0098	0.0261	0.0417	0.1228
		(0.1117)	(0.0202)	(0.0119)	(0.0042)	(0.0028)
	β_1	0.0387	0.0822	0.0851	0.0843	0.1663
		(0.0000)	(0.0000)	(0.0000)	(0.0000)	(0.0000)
	β_2	0.9586	0.9149	0.9021	0.8928	0.8023
		(0.0000)	(0.0000)	(0.0000)	(0.0000)	(0.0000)
Skewness equation	γ_0	-0.0458	-0.0886	-0.0245	-0.0446	0.0228
		(0.0518)	(0.0106)	(0.2911)	(0.0161)	(0.3101)
	γ_1	0.0085	0.0078	0.0048	0.0189	0.0125
		(0.0139)	(0.0032)	(0.2006)	(0.0000)	(0.0136)
	γ_2	0.0227	0.2174	0.6781	0.1352	0.2969
		(0.9187)	(0.4136)	(0.0168)	(0.0852)	(0.3112)
Kurtosis equation	δ_0	3.0471	1.4576	0.4866	0.2526	0.3302
		(0.0000)	(0.0175)	(0.0016)	(0.0026)	(0.0254)
	δ_1	0.0055	0.0007	0.0010	0.0004	0.0010
		(0.0019)	(0.6228)	(0.0229)	(0.0129)	(0.3634)
	δ_2	0.0882	0.5518	0.8493	0.9208	0.9018
		(0.5715)	(0.0034)	(0.0000)	(0.0000)	(0.0000)
Log-Likelihood	-	-1404.5752	-2375.0218	-2484.1335	-2414.6928	-2056.0966
SIC	-	-1445.2519	-2415.7000	-2525.7985	-2455.3328	-2094.4277

TABLE 8
NAGARCHSK MODELS – STOCK INDICES

The reported coefficients shown in each row of the table are ML estimates of the NAGARCHSK model:

$$r_t = \alpha_1 r_{t-1} + \varepsilon_t$$

$$h_t = \beta_0 + \beta_1 (\varepsilon_{t-1} + \beta_3 h_{t-1}^{1/2})^2 + \beta_2 h_{t-1}$$

$$s_t = \gamma_0 + \gamma_1 \eta_{t-1}^3 + \gamma_2 s_{t-1}$$

$$k_t = \delta_0 + \delta_1 \eta_{t-1}^4 + \delta_2 k_{t-1}$$

for percentage daily returns of S&P500, NASDAQ100, DAX30, IBEX35 stock indices, from January 1990 to July 2003, and MEXBOL from January 1995 to July 2003. $h_t = \text{var}(r_t | r_{t-1}, r_{t-2}, \dots)$, $s_t = \text{skewness}(r_t | r_{t-1}, r_{t-2}, \dots)$, $k_t = \text{kurtosis}(r_t | r_{t-1}, r_{t-2}, \dots)$, $\eta_t = \varepsilon_t h_t^{-1/2}$, and $\varepsilon_t | \varepsilon_{t-1}, \varepsilon_{t-2}, \dots$ follows the distribution based on a Gram-Charlier series expansion. All models have been estimated by ML using the Berndt-Hall-Hall-Hausman algorithm (quasi-maximum likelihood p-values in parenthesis; in bold are significantly different from zero coefficients).

	Parameter	S&P500	NASDAQ	DAX30	IBEX35	MEXBOL
Mean equation	α_1	0.0358 (0.0466)	0.1255 (0.0000)	0.0152 (0.4009)	0.1024 (0.0000)	0.1742 (0.0000)
	Variance equation	β_0	0.0083 (0.0006)	0.01841 (0.0038)	0.0278 (0.0005)	0.04460 (0.0004)
β_1		0.0416 (0.0000)	0.0986 (0.0000)	0.0696 (0.0000)	0.0729 (0.0000)	0.1202 (0.0000)
β_2		0.9099 (0.0373)	0.8801 (0.0000)	0.8961 (0.0000)	0.8800 (0.0000)	0.7834 (0.0000)
β_3		-1.0116 (0.0000)	-0.4351 (0.0000)	-0.5597 (0.0000)	-0.5795 (0.0003)	-0.7703 (0.0000)
Skewness equation	γ_0	-0.0451 (0.0373)	-0.0618 (0.0005)	-0.0261 (0.2285)	-0.0204 (0.1174)	0.0525 (0.0782)
	γ_1	0.0091 (0.1034)	0.0103 (0.0025)	0.0050 (0.1883)	0.0045 (0.1423)	0.0180 (0.0045)
	γ_2	0.0552 (0.7418)	0.4572 (0.0000)	0.6573 (0.0124)	0.5325 (0.0022)	0.1922 (0.5459)
Kurtosis equation	δ_0	3.1652 (0.0000)	1.6929 (0.0003)	0.4536 (0.0016)	0.2012 (0.0858)	1.9901 (0.0011)
	δ_1	0.0150 (0.0000)	0.0053 (0.0025)	0.0009 (0.0161)	0.0004 (0.0749)	0.0055 (0.0004)
	δ_2	0.0293 (0.6645)	0.4684 (0.0014)	0.8581 (0.0000)	0.9365 (0.0000)	0.4017 (0.0271)
Log-Likelihood	-	-1371.4169	-2351.1665	-2461.0251	-2382.5437	-2016.8569
SIC	-	-1416.1613	-2395.9126	-2505.7566	-2427.2477	-2059.0212

TABLE 9
LIKELIHOOD RATIO TESTS

The table shows the values of the maximized log-likelihood function (logL) when the distribution for the error term is assumed to be normal (standard GARCH or NAGARCH specification) and when it is assumed to be a Gram-Charlier series expansion of the normal density (GARCHSK or NAGARCHSK specification), the likelihood ratio (LR) and asymptotic p-values for the series employed in the paper (in bold are significantly different from zero LR statistics)

PANEL A: EXCHANGE RATES					
STATISTIC	GBP/USD	JPY/USD	DEM/USD	CHF/USD	
LogL(GARCH)	409.3	-352.6	-149.3	-451.7	
LogL(GARCHSK)	472.4	-237.7	-117.6	-421.0	
LR	126.1	229.9	63.4	61.5	
(p-value)	(0.00)	(0.00)	(0.00)	(0.00)	

PANEL B: STOCK INDICES					
STATISTIC	S&P500	NASDAQ100	DAX30	IBEX35	MEXBOL
LogL(NAGARCH)	-1401.9	-2385.4	-2496.0	-2413.7	-2050.1
LogL(NAGARCHSK)	-1371.4	-2351.2	-2461.0	-2382.5	-2016.9
LR	60.9	68.4	70.0	62.3	72.8
(p-value)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)

TABLE 10
DESCRIPTIVE STATISTICS FOR CONDITIONAL VARIANCES

The table shows the main descriptive statistics for the conditional variances obtained from GARCH and GARCHSK models for GBP/USD series, and from NAGARCH and NAGARCHSK models for S&P500 series paper (in bold are significantly different from zero Jarque-Bera statistics)

STATISTIC	GBP/USD		S&P500	
	h_t - GARCH	h_t - GARCHSK	h_t - NAGARCH	h_t - NAGARCHSK
Simple size	3124	3124	3413	3413
Mean	0.3264	0.3026	1.1394	1.0928
Median	0.2647	0.2432	0.7692	0.7513
Maximum	1.4762	1.3944	8.3534	6.9340
Minimum	0.0988	0.0776	0.1731	0.1771
Stand. Dev.	0.2034	0.1980	1.0575	0.9533
Skewness	2.2384	2.1624	2.5160	2.2077
Kurtosis	9.4659	8.9007	11.1431	8.9475
Jarque-Bera (p-value)	8050.721 (0.0000)	6966.893 (0.0000)	13030.790 (0.0000)	7802.598 (0.0000)

TABLE 11
IN-SAMPLE PREDICTIVE POWER

The variable predicted is the squared forecast error (ε_t^2) and the predictors are the conditional variances (h_t) from, respectively, the standard GARCH or NAGARCH models and GARCHSK or NAGARCHSK models. Two metrics are chosen to compare the predictive power ability of these models:

1. Median absolute error $MAE = med(|\varepsilon_t^2 - h_t|)$
2. Median percentage absolute error $MPAE = med\left(\frac{|\varepsilon_t^2 - h_t|}{\varepsilon_t^2}\right)$

The metrics are based on the median given the high dispersion of the error series.

SERIES		MAE	MPAE
GBP/USD	G	0.2030	1.9227
	GSK	0.1874	1.6567
JPY/USD	G	0.3369	2.2226
	GSK	0.3165	2.0134
DEM/USD	G	0.3058	1.7982
	GSK	0.2895	1.6028
CHF/USD	G	0.3749	1.8096
	GSK	0.3635	1.6788
S&P500	NG	0.5884	1.7690
	NGSK	0.5723	1.7670
NASDAQ	NG	0.9061	1.3801
	NGSK	0.9209	1.3075
DAX30	NG	1.0225	1.5102
	NGSK	1.0207	1.5071
IBEX35	NG	1.0081	1.4610
	NGSK	1.0109	1.4349
MEXBOL	NG	1.6743	1.6508
	NGSK	1.6308	1.5531