

Parental education and genetics of body mass index from infancy to old age:
a pooled analysis of 29 twin cohorts

Karri Silventoinen (1) (2), Aline Jelenkovic (1) (3), Antti Latvala, (4) (5), Yoshie Yokoyama (6), Reijo Sund (1) (7), Masumi Sugawara (8), Mami Tanaka (9), Satoko Matsumoto (10), Sari Aaltonen (1) (5), Maarit Piirtola (1) (5), Duarte L Freitas (11), José A Maia (12), Sevgi Y Öncel (13), Fazil Aliev (14), Fuling Ji (15), Feng Ning (15), Zengchang Pang (15), Esther Rebato (3), Kimberly J Saudino (16), Tessa L Cutler (17), John L Hopper (17) (18), Vilhelmina Ullemar (19), Catarina Almqvist (19) (20), Patrik KE Magnusson (19), Wendy Cozen (21) (22), Amie E Hwang (21), Thomas M Mack (21) (22), Gonneke Willemsen (23), Meike Bartels (23), Catharina EM van Beijsterveldt (23), Tracy L Nelson (24), Keith E Whitfield (25), Joohon Sung (18) (26), Jina Kim (18), Jooyeon Lee (18), Sooji Lee (18), Clare H Llewellyn (27), Abigail Fisher (27), Emanuela Medda (28), Lorenza Nisticò (28), Virgilia Toccaceli (28), Laura A Baker (29), Catherine Tuvblad (29) (30), Robin P Corley (31), Brooke M Huibregtse (32), Catherine A Derom (33) (34), Robert F Vlietinck (33), Ruth JF Loos (35), Ariel Knafo-Noam (36), David Mankuta (37), Lior Abramson (36), S Alexandra Burt (38), Kelly L Klump (38), Judy L Silberg (39), Hermine H Maes (40), Robert F Krueger (41), Matt McGue (41), Shandell Pahlen (41), Margaret Gatz (19) (42), David A Butler (43), Jennifer R Harris (44), Thomas S Nilsen (44), K Paige Harden (45), Elliot M Tucker-Drob (45), Carol E Franz (46), William S Kremen (46) (47), Michael J Lyons (48), Paul Lichtenstein (19), Hoe-Uk Jeong (49), Yoon-Mi Hur (49), Dorret I Boomsma (23), Thorkild IA Sørensen (50) (51), Jaakko Kaprio (4) (5)

1. Department of Social Research, University of Helsinki, Helsinki, Finland.
2. Osaka University Graduate School of Medicine, Osaka University, Osaka, Japan.
3. Department of Genetics, Physical Anthropology and Animal Physiology, University of the Basque Country UPV/EHU, Leioa, Spain.
4. Department of Public Health, University of Helsinki, Helsinki, Finland.
5. Institute for Molecular Medicine Finland (FIMM), Helsinki, Finland.
6. Department of Public Health Nursing, Osaka City University, Osaka, Japan.
7. Institute of Clinical Medicine, University of Eastern Finland, Kuopio, Finland.
8. Department of Psychology, Ochanomizu University, Tokyo, Japan.
9. Center for Forensic Mental Health, Chiba University, Chiba, Japan.
10. Institute for Education and Human Development, Ochanomizu University, Tokyo.
11. Department of Physical Education and Sport, University of Madeira, Funchal, Portugal.
12. CIFI2D, Faculty of Sport, Porto, University of Porto, Portugal.
13. Department of Statistics, Faculty of Arts and Sciences, Kırıkkale University, Kırıkkale, Turkey.
14. Psychology and African American Studies, Virginia Commonwealth University, Richmond, VA, USA.
15. Department of Noncommunicable Diseases Prevention, Qingdao Centers for Disease Control and Prevention, Qingdao, China.
16. Boston University, Department of Psychological and Brain Sciences, Boston, MA, USA.
17. The Australian Twin Registry, Centre for Epidemiology and Biostatistics, University of Melbourne, Melbourne, Victoria, Australia.
18. Department of Epidemiology, School of Public Health, Seoul National University, Seoul, Korea.
19. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden.
20. Pediatric Allergy and Pulmonology Unit at Astrid Lindgren Children's Hospital, Karolinska University Hospital, Stockholm, Sweden.

21. Department of Preventive Medicine, Keck School of Medicine of USC, University of Southern California, Los Angeles, California, USA.
22. USC Norris Comprehensive Cancer Center, Los Angeles, California, USA.
23. Netherlands Twin Register, Department of Biological Psychology, Vrije Universiteit Amsterdam, Amsterdam, Netherlands..
24. Department of Health and Exercise Sciences and Colorado School of Public Health, Colorado State University, Fort Collins, Colorado, USA.
25. Psychology and Neuroscience, Duke University, Durham, NC, USA.
26. Institute of Health and Environment, Seoul National University, Seoul, South-Korea.
27. Health Behaviour Research Centre, Department of Epidemiology and Public Health, Institute of Epidemiology and Health Care, University College London, London, UK.
28. Centre for Behavioural Sciences and Mental Health, Istituto Superiore di Sanità - Rome, Italy.
29. Department of Psychology, University of Southern California, Los Angeles, CA, USA.
30. School of Law, Psychology and Social Work, Örebro University, Örebro, Sweden.
31. Institute for Behavioral Genetics, University of Colorado, Boulder, Colorado, USA.
32. Institute of Behavioral Science, University of Colorado, Boulder, Colorado, USA.
33. Centre of Human Genetics, University Hospitals Leuven, Leuven, Belgium.
34. Department of Obstetrics and Gynaecology, Ghent University Hospitals, Ghent, Belgium.
35. The Charles Bronfman Institute for Personalized Medicine, The Mindich Child Health and Development Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA.
36. The Hebrew University of Jerusalem, Jerusalem, Israel.
37. Hadassah Hospital Obstetrics and Gynecology Department, Hebrew University Medical School, Jerusalem, Israel.
38. Michigan State University, East Lansing, Michigan, USA.
39. Department of Human and Molecular Genetics, Virginia Institute for Psychiatric and Behavioral Genetics, Virginia Commonwealth University, Richmond, Virginia, USA.
40. Department of Human and Molecular Genetics, Psychiatry & Massey Cancer Center, Virginia Commonwealth University, Richmond, Virginia, USA.
41. Department of Psychology, University of Minnesota, Minneapolis, MN, USA.
42. Center for Economic and Social Research, University of Southern California, Los Angeles, CA, USA.
43. Health and Medicine Division, The National Academies of Sciences, Engineering, and Medicine Washington, DC, USA.
44. Norwegian Institute of Public Health, Oslo, Norway.
45. Department of Psychology, University of Texas at Austin, Austin, TX, USA.
46. Department of Psychiatry, University of California, San Diego, CA, USA.
47. VA San Diego Center of Excellence for Stress and Mental Health, La Jolla, CA, USA.
48. Boston University, Department of Psychology, Boston, MA, USA.
49. Department of Education, Mokpo National University, Jeonnam, South Korea.
50. Novo Nordisk Foundation Centre for Basic Metabolic Research (Section of Metabolic Genetics), Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.
51. Department of Public Health (Section of Epidemiology), Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.

Key words: body mass index, parental education, genetics, twins, international comparisons

Conflict of interests: The authors declare no competing financial interests.

Running title: Parental education and genetics of BMI

Contact information:

Karri Silventoinen

University of Helsinki, Department of Social Research

P.O. Box 18, FIN-00014 University of Helsinki, Finland

GSM: +358400-620726

E-mail: karri.silventoinen@helsinki.fi

Word count: 3996 words

Funding: This study was conducted within the CODATwins project (Academy of Finland #266592). Funding for individual collaborators: The CATSS-Study is supported by the Swedish Research Council through the Swedish Initiative for Research on Microdata in the Social And Medical Sciences (SIMSAM) framework grant no 340-2013-5867, grants provided by the Stockholm County Council (ALF-projects), the Swedish Heart-Lung Foundation and the Swedish Asthma and Allergy Association's Research Foundation. Netherlands Twin Register acknowledges the Netherlands Organization for Scientific Research (NWO) and MagW/ZonMW grants 904-61-090, 985-10-002, 912-10-020, 904-61-193,480-04-004, 463-06-001, 451-04-034, 400-05-717, Addiction-31160008, Middelgroot-911-09-032, Spinozapremie 56-464-14192; VU University's Institute for Health and Care Research (EMGO+); the European Research Council (ERC - 230374), the Avera Institute, Sioux Falls, South Dakota (USA). Data collection and analyses in Finnish twin cohorts have been supported by ENGAGE – European Network for Genetic and Genomic Epidemiology, FP7-HEALTH-F4-2007, grant agreement number 201413, National Institute of Alcohol Abuse and Alcoholism (grants AA-12502, AA-00145, and AA-09203 to R J Rose, the Academy of Finland Center of Excellence in Complex Disease Genetics (grant numbers: 213506, 129680), and the Academy of Finland (grants 100499, 205585, 118555, 141054, 265240, 264146 and 312073 to J Kaprio). Since its origin the East Flanders Prospective Survey has been partly supported by grants from the Fund of Scientific Research, Flanders and Twins, a non-profit Association for Scientific Research in Multiple Births (Belgium). Gemini was supported by a grant from Cancer Research UK (C1418/A7974). Data collection and research stemming from the Norwegian Twin Registry is supported, in part, from the European Union's Seventh Framework Programmes ENGAGE Consortium (grant agreement HEALTH-F4-2007-201413, and BioSHaRE EU (grant agreement HEALTH-F4-2010-261433). Madeira data comes from the following project: Genetic and environmental influences on physical activity, fitness and health: the Madeira family study Project reference: POCI/DES/56834/2004 Founded by the Portuguese agency for research (The Foundation for Science and Technology [FCT]). S.Y. Öncel and F. Aliev are supported by Kırıkkale University Research Grant: KKKU, 2009/43 and TUBITAK grant 114C117. K Silventoinen is supported by Osaka University's International Joint Research Promotion Program. The Boston University Twin Project is funded by grants (#R01 HD068435 #R01 MH062375) from the National Institutes of Health to K. Saudino. California Twin Program was supported by The California Tobacco-Related Disease Research Program (7RT-0134H, 8RT-0107H, 6RT-0354H) and the National Institutes of Health (1R01ESO15150-01). The Carolina African American Twin Study of Aging (CAATSA) was funded by a grant from the National Institute on Aging (grant 1R01-AG13662-01A2) to K. E. Whitfield. Colorado Twin Registry is funded by NIDA funded center grant DA011015, &

Longitudinal Twin Study HD10333; Author Huibregtse is supported by 5T32DA017637 and 5T32AG052371. The Michigan State University Twin Registry has been supported by Michigan State University, as well as grants R01-MH081813, R01-MH0820-54, R01-MH092377-02, R21-MH070542-01, R03-MH63851-01 from the National Institute of Mental Health (NIMH), R01-HD066040 from the Eunice Kennedy Shriver National Institute for Child Health and Human Development (NICHD), and 11-SPG-2518 from the MSU Foundation. The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the NIMH, the NICHD, or the National Institutes of Health. The University of Southern California Twin Study is funded by a grant from the National Institute of Mental Health (R01 MH58354). The Texas Twin Project is currently funded by grants AA023322 and HD081437 from the National Institutes of Health. Vietnam Era Twin Study of Aging was supported by National Institute of Health grants NIA R01 AG018384, R01 AG018386, R01 AG022381, and R01 AG022982, and, in part, with resources of the VA San Diego Center of Excellence for Stress and Mental Health. The Cooperative Studies Program of the Office of Research & Development of the United States Department of Veterans Affairs has provided financial support for the development and maintenance of the Vietnam Era Twin (VET) Registry. The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the NIA/NIH, or the VA. The NAS-NRC Twin Registry acknowledges financial support from the National Institutes of Health grant number R21 AG039572. Korean Twin-Family Register was supported by the Global Research Network Program of the National Research Foundation (NRF 2011-220-E00006). South Korea Twin Registry is supported by National Research Foundation of Korea (NRF-371-2011-1 B00047). The West Japan Twins and Higher Order Multiple Births Registry was supported by Grant-in-Aid for Scientific Research (B) (grant number 15H05105) from the Japan Society for the Promotion of Science. This research was facilitated through access to Twins Research Australia, a national resource supported by a Centre of Research Excellence Grant (ID: 1079102), from the National Health and Medical Research Council. Longitudinal Israeli Study of Twins was funded by the Starting Grant no. 240994 from the European Research Council (ERC) to Ariel Knafo.

Disclosure: J Kaprio reports grants from Academy of Finland during the conduct of the study.

What is already known about this subject?

- 1) Low parental education is associated with greater body mass index in offspring from early childhood until adulthood.
- 2) Genetic factors explain a large proportion of the variation of body mass index, and there is evidence that environmental factors can modify these effects.

What this study adds?

- 1) Using pooled data from 29 twin cohorts, we showed that low parental education is associated with greater genetic variation of body mass index from early childhood to old age.
- 2) The interaction effect between parental education and genetic factors on body mass index was stronger in North America and Australia than in Europe and East Asia.
- 3) Genetic factors intermediate the effect of childhood social position on body mass index seen over the human life course, and this association is strongest in the obesogenic social context.

Abstract

Objective: To analyze how parental education modifies the genetic and environmental variances of body mass index (BMI) from infancy to old age in three geographic-cultural regions.

Methods: A pooled sample of 29 cohorts including 143 499 twin individuals with information on parental education and BMI from 1 to 79 years (299 201 BMI measures) was analyzed by genetic twin modeling.

Results: Until 4 years of age, parental education was not consistently associated with BMI. Thereafter, higher parental education was associated with lower BMI in males and females. Total and additive genetic variances of BMI were smaller in the offspring of highly educated parents than in those whose parents had low education. Especially in North American and Australian children, environmental factors shared by co-twins also contributed to the higher BMI variation in the low educational category. In Europe and East Asia, the associations of parental education with mean BMI and BMI variance were weaker than in North America and Australia.

Conclusions: Lower parental education is associated with higher mean and larger genetic variance of BMI after early childhood, especially in the obesogenic macro-environment. The interplay between genetic predisposition, childhood social environment and macro-social context is important for socio-economic differences in BMI.

Introduction

Mean body mass index (BMI, kg/m²) has increased all over the industrialized world during the last four decades and currently shows clear geographic differences with the highest BMIs in North America and lowest in East Asia (1). Twin studies have shown that genetic factors explain a substantial proportion of BMI variation over the life course (2,3), and genome wide association (GWA) studies have identified the role of genetic variants behind BMI variation (4,5). There is also strong evidence for a higher BMI in lower socio-economic classes in Western industrialized societies (6). The roots of socio-economic obesity differences may already emerge in childhood since many socio-economic characteristics of childhood family, such as parental education and income, are associated with the BMI of offspring (7). Adoption studies supported these findings by showing that lower socio-economic status of biologically unrelated adoptive families is associated with higher BMI of adopted children as adults (8).

The association of genes and environmental factors with BMI may be modified by socio-demographic characteristics. In our previous study based on the same database as in this present study, we found that the genetic variation of adult BMI increased from the 1940s to the 2010s at the same time when mean BMI dramatically increased (3) confirming the earlier findings based on Danish (9) and Swedish data (10). A Dutch twin study of parental education in children and adolescents (11) as well as studies on Danish (12) and US adults (13) of their own education suggested increased genetic BMI variation associated with lower education. However, a Finnish twin study reported that in adolescence the increased BMI variation in the offspring of parents with lower education was attributable to environmental factors shared by co-twins (14).

To address the question how modification of genetic and environmental influences on BMI develops over the human lifespan requires considerable statistical power. Thus, we analyzed if and how parental education modifies the genetic and environmental variation of BMI from infancy to old age in a very large international database of twins. Further, we analyzed whether the interaction between parental education and BMI varies between three geographic-cultural regions that are characterized by different levels of mean BMI in the population indicating different obesogenic environments.

Methods

The data were derived from the CODATwins (COllaborative project of Development of Anthropometrical measures in Twins) project targeted to collect all available twin data on height and weight in the world (15). We included 29 twin cohorts with additional information on maternal and paternal education. Eleven cohorts came from European countries, five from East Asian countries, eleven from the USA, and single cohorts from Australia and Israel. The names of participating cohorts are given in Table 1 (footnote). Age was classified to one-year age categories from 1 to 19 years of age, and after that 10-year categories were defined. Participants who were 80 years of age or older were excluded (319 measures) because of decreasing muscle mass at old age (16). Together, there were 324 329 BMI measurements. In children and adolescents, we checked the BMI distribution in each age and sex group and removed the extreme, likely erroneous values (658 measures representing 0.3% of the measurements at 19 years of age or younger corresponding to values approximately below or above 3 SD from the mean). In adults, we removed the measurements consistent with anorexia nervosa (BMI<15 kg/m², 96 measures representing 0.1% of adult measures) and morbid obesity (BMI>45 kg/m², 246 measures representing 0.3% of adult measures). Thus, our results represent the range of normal BMI variation, and not extremely low and very high BMI values, which may be affected

specific factors such as anorexia nervosa or rare monogenic traits. To confirm that all parameter estimates were based on independent observations, we selected one measurement in each age group from cohorts with repeated measures and conducted independent models in each age group. In the present analyses, 299 201 BMI measures from 143 499 twin individuals (52% females) including 66 165 complete twin pairs (39% monozygotic (MZ), 34% same-sex dizygotic (DZ) and 27% opposite-sex DZ) with information on maternal and paternal education were used. All participants were volunteers who gave informed consent when participating in their original studies. The pooled analysis was approved by the ethical committee of the Department of Public Health, University of Helsinki.

The different educational classifications used in the surveys were transformed as educational years as described elsewhere (17). We first calculated regression residuals of maternal and paternal education separately by fitting a regression model with maternal or paternal education as the dependent variable and twin cohort and the birth year of their twin children, used as a proxy indicator for the birth years of parents, as independent variables. Twin cohort was treated as a nominal level variable with a dummy variable for each individual cohort and birth year as a continuous variable. These residuals were then summed up to get combined parental education and divided into three categories (less than -0.5 SD from the mean, from -0.5 until 0.5 SD from the mean and more than 0.5 SD from the mean) to indicate low (31% of participants), average (40% of participants) and high parental education (29% of participants). The SD of parental education was slightly higher in North America and Australia (SD=5.33) than in Europe (SD=4.50) and East Asia (SD=4.10). However, we calculated the residuals in the full cohort and used the same categorization of parental education to avoid changing the educational distribution and to confirm the same distance of education in all geographic-cultural regions. Thus, the residuals indicate how much shorter or longer the parental education is as compared

to the average person having a certain birth year in each twin cohort. Correspondingly, BMI was adjusted for the effects of exact age and twin cohort in each age and sex category by regression models.

In the aggregation of twin cohorts by the geographic-cultural regions, we removed Turkey from all region specific analyses since the Turkish population is genetically distinct from other European populations (18). We also combined the cohort from Australia with North American cohorts since the mean BMI in Australia is closer to North American than European countries (1). Thus, we had three geographic-cultural regions, North America and Australia, Europe, and East Asia representing the high, intermediate and low level of obesogenic environment, respectively, based on the mean BMI of population (1). The same classification was used also in our previous studies on the genetics of BMI in childhood (2) and adulthood (3) based on the CODATwins database. The number of BMI measurements was highest in Europe (196 990 measures) followed by North America and Australia (82 098 measures) and East Asia (18 670 measures). Since there was more ethnic diversity in North America and Australia than in the other cultural-geographic regions, we repeated the main analyses in North Americans and Australians of Caucasian ethnicity based on self-reports (75 973 measures).

We used structural equation modeling to estimate genetic and environmental variance components (19). Genetic twin modeling is based on the different genetic relatedness of MZ and DZ twins: DZ twins share, on average, 50% of their genes identical by descent; MZ twins are virtually identical at the DNA sequence level. Based on this principle, the BMI variation can be decomposed into additive genetic variation (additive effects of all loci affecting BMI), dominance genetic variation (non-additive genetic effects), shared environmental variation (all environmental effects making co-twins similar) and unique environmental variation (all environmental effects making co-twins different and measurement error). Models were fitted separately for each parental education category by age and sex

groups. As we reported previously, slightly higher variances of BMI were found from 6 to 12 years of age in DZ than in MZ twins (20). However, these differences were minor, i.e. 15% of less, and became statistically significant only because of our very large sample size.

As we have previously reported for children (2) and adults (3), there was little evidence for the presence of dominance genetic variation in BMI. Thus, we used the additive genetic/ shared environment/ unique environment model in the analyses, which was also supported by the within pair correlations in all parental education categories in our study cohort (Table S1). We fitted the full model in each age group to get the estimation of the role of shared environmental factors, even when it was not statistically significant in most of age categories. A clear sex-specific genetic effect for BMI was found in childhood (2) and adulthood (3), and thus all models were fitted allowing the opposite-sex DZ genetic correlation to be lower than the 0.5 expected for same-sex DZ twins. Especially in early childhood, DZ twins were slightly heavier than MZ twins (20), and thus a different mean BMI was used for MZ and DZ twins.

Next, we fitted gene-environment interaction models to test the significance of the moderator effect of parental education (21). In these models, parental education, used as a continuous variable, was allowed to modify the genetic and environmental variation of BMI. Since weak but systematic shared environmental effects were found for BMI in this database, especially in childhood (2), we fitted both a more parsimonious additive genetic/ unique environment (AE) model and an additive genetic/ shared environment/ unique environment (ACE) model, which needs more statistical power. The genetic models were fitted to the data by the OpenMx package (version 2.0.1) of R statistical software (22). The effects of parental education on mean BMI were estimated by Stata/SE 13.1 for Windows statistical software. The effect of intra-pair correlations on 95% confidence intervals (CI) was taken

into account by the cluster option of Stata. Further, we adjusted these mean modification effects for zygosity because of slight differences in BMI (20) and parental education between MZ and DZ twins (17).

Results

Table 1 presents the descriptive statistics by age and sex. Mean BMI shows the nadir at 5 years of age and the steady increase until 50-59 years of age with modest sex differences. Mean BMI was highest in North America and Australia and lowest in East Asia (Table S2). Maternal and paternal education decreased from the younger to the older age groups reflecting the increasing education over the birth cohorts. As expected, parental education was virtually identical for males and females.

Figure 1 presents the associations between parental education and BMI. No inverse educational association was seen from 1 to 3 years of age, and the sons of better-educated parents had even a slightly higher BMI. An inverse educational association emerged at 5 years of age in boys and 4 years of age in girls and became stronger until adolescence. In adulthood, the inverse association between parental education and BMI was statistically significant in most of the age groups and stronger in women than in men. In men this association was somewhat weaker in late adulthood as compared with middle-age, but otherwise no systematic differences between the age groups in adulthood were seen.

Figure 2 presents the associations between parental education and BMI in the three cultural-geographic regions. Since most of the BMI measures came from European cohorts, the pattern in Europe followed closely the estimates in the pooled cohort. In North America and Australia, parental education showed stronger associations with mean BMI than in Europe in most of age groups. These differences were

more pronounced in females than in males. In East Asia, similar associations were found in females, but because of wider 95% CIs, they were not statistically significant. However, East Asian males showed no evidence on the association between parental education and BMI.

The total variation of BMI in the three categories of parental education, decomposed to additive genetic, shared environmental and unique environmental variances, is shown in Figure 3 (the exact estimates with 95% CIs are provided in Table S3). The total BMI variation was greatest in the low parental education category and smallest in the high parental education category in all age groups. With a few exceptions, the BMI variation in the intermediate parental education category was between the high and low education categories. In nearly all age groups, the difference in BMI variation between the low and high parental education categories was statistically significant as seen by non-overlapping 95% CIs (Table S3). When we studied the components of variation, a similar pattern was observed with a few exceptions for additive genetic and unique environmental variances. Except in two age groups, the difference was statistically significant for additive genetic variation, but for unique environmental variation, in nearly half of the age groups the 95% CIs overlapped. Shared environmental variance components explained part of the variance, especially in boys and girls of 14 years of age or younger. However, in most of the age groups, they were not statistically significant, and the 95% CIs also overlapped between the educational categories (Table S3).

The gene-environment interaction models confirmed the results of the stratified analyses (Table S4). When we fitted the AE model, in all age groups except 70-79 years old men, additive genetic modification effects were negative, i.e. higher parental education was associated with less genetic variation. However, in men 17 years of age or older and in girls three years or younger and women 50-59 years or older, the modification effects were weaker than in the other age groups and not significant.

Most of unique environmental modification effects were also negative, but weaker than additive genetic modification effects, and around half of them were not statistically significant. In the ACE model, most of the additive genetic and shared environmental modification effects were negative, but there was more fluctuation in the estimates because of lesser statistical power, and only part of them were statistically significant.

Figures 4 and 5 depict the similar decomposition of BMI variance by the cultural-geographic region. Since the results for the intermediate parental educational category were, in general, between the high and low educational categories, they are not shown, but all estimates with 95% CIs are available in Table S5 and Table S6. The total BMI variance was greatest in North America and Australia and smallest in East Asia. In Europe, as well as in North America and Australia, BMI variation was systematically greater in males and females whose parents have low education as compared to the offspring of highly educated parents. In most of the age groups, this difference was statistically significant as seen as non-overlapping 95% CIs (Tables S5 and Table S6). When we restricted the analyses to North Americans and Australians with Caucasian ethnicity, the variances decreased somewhat, but they were still larger than in the other cultural-geographic regions and highest in those with low parental education (Table S5 and Table S6).

The differences in the total BMI variation between the parental educational categories were mainly due to differences in additive genetic variation. An exception was North American and Australian boys and girls 11 years of age or younger, for whom the difference in most age groups was mainly due to shared environmental variation. In Europe, we also observed some evidence for the greater role of shared environment in the low parental education category until 14 years of age, but the differences were not

statistically significant. In East Asia, no systematic differences were seen in the total variance of BMI according to parental education.

The gene-environment interaction models confirmed these results (Table S7). In Europe, the additive genetic modification effects were consistently negative, whereas there was more fluctuation in the common environmental modifications effects. In contrast, in North America and Australia, especially in boys under 12 years of age, the shared environmental modifications were consistently negative. However when using the ACE models, which require more statistical power, only part of these modifications effects were statistically significant. In East Asia, the modification effects fluctuated in both sides of zero and most of them were not statistically significant.

Discussion

In this study of a very large international dataset of twins, we found that higher parental education was consistently associated with lower additive genetic and unique environmental variance of BMI. This result is in accordance with previous twin and molecular genetic studies. In a previous study based on Dutch twin data, children and adolescents whose parents had middle or low education showed greater genetic BMI variance than those whose parents had high education (3). Danish (12) and US (13) studies of adult twins supported this result by showing greater genetic variation in those with lower than in those with higher education. These results are also in line with candidate genes studies in which low maternal education in children (23) and own education in adults (24) reinforced the effect of FTO gene, the major candidate gene of BMI. Similar results were found for other BMI candidate genes (25,26). However, these molecular genetic studies need replications in other populations, and preferably a GWA study using a large sample size should be conducted. Assuming parental education

to be a good proxy indicator of the socio-economic conditions of individuals and their families, our results might indicate that poorer childhood socio-economic environment reinforces the influence of genetic factors affecting BMI. However, parental education is associated also with own education in adulthood (27), which is further associated with other factors such as intelligence and personality (28). Thus, the mediating pathways are complex and may change through the life-course.

Parental education is associated with BMI after early childhood, which parallels previous population based studies showing no association (29) or lower BMI (30) in the offspring of mothers with lower education before two years of age but a clear inverse correlation between the BMI of offspring and maternal education after four years of age. In early childhood, the lack of association between parental education and offspring BMI might be due to low birth weight associated with low parental education affecting BMI over the first years (29). Nevertheless, it may also reflect a change in the genetic architecture of BMI. A negative genetic correlation between education and BMI ($r=-0.28$, standard error=0.03) was found based on cross-trait linkage-disequilibrium score regression analysis of large GWA studies (31). As we have reported previously, genetic variation in BMI started to increase after 5 years of age in the CODATwins database (2). This result is in accordance with the previous molecular genetic studies showing that the effects of the FTO gene (32), genetic risk scores of top BMI hits (33,34) and polygenic risk scores from GWA studies (35) started to increase only after early childhood. Thus, it seems that after early childhood, new genetic variants affecting BMI activate and parental education can modify the effects of them, which may be independent of the genetic variants regulating body size in infancy.

The mechanisms explaining the interaction effects between genes and socio-economic factors on BMI are not clear. In the present study, the association of parental education with mean and genetic variance

of BMI persisted through adulthood. This trend might be due to behavior. There is evidence that the FTO gene may affect BMI through food-intake self-regulation (36) and eating styles (37), and many genetic risk variants of adult BMI are active in brain areas having an important role in appetite regulation and many cognitive functions (4,5). Findings that the correlations between BMI and a number of personality, cognitive and brain morphological traits were largely explained by common genetic factors further support the mediating role of brain function (38). It is thus possible that the new genetic variance emerging after mid-childhood relates to the child's food-intake self-regulation, with children learning better self-regulation in households with higher parental education. Childhood social position also relates to factors such as diet, adverse childhood experiences and physical activity (39). There is evidence that physical activity can modify the association between the BMI genetic risk-score and BMI (40), and parallel results have been found in a twin study showing greater genetic variance of BMI and other obesity indicators in sedentary persons as compared to those who are physically active (41). However, the background mechanisms may go even deeper in the field of neuro-physiology. Subordinate female rhesus monkeys had higher caloric consumption than dominant monkeys (42). Also starlings with a disadvantaged early life environment were fatter in adulthood and spent more effort for food gathering than those with more advantaged early life environments (43). Although comparisons between species should be treated with caution, these results may suggest deep neurophysiological pathways between early life social position and later eating behavior.

When comparing the geographic-cultural regions, parental education was more strongly associated with BMI in North America and Australia than in Europe and East Asia. Thus, obesogenic environment may reinforce the association between BMI and parental education since the population level mean of BMI is much higher in North America and Australia than in the other two regions (1). This would suggest that the factors affecting BMI at population level have stronger effects on those having social and/or

genetic susceptibility to gain weight. There has been discussion regarding to which specific societal factors are behind these regional level differences in BMI, but the results are inconclusive (44). However, our findings suggest that identifying socio-economic factors associated with the increase of BMI would have beneficial effects especially on those who are most vulnerable to gain weight. Also other societal factors than obesogenic environment may lead to the weaker effect of parental education on BMI in East Asia. For example, because of cultural differences, food consumption may be differently associated with social factors in different cultural-geographic regions even on the same level of obesogenic environment. More detailed measures of macro-environment would be needed to further disentangle the effect of these societal factors.

In most of the age categories, the differences in BMI variance were mainly due to the greater additive genetic variance in the lower parental education category. However, especially in North America and Australia, this difference was due to greater shared environmental variance in boys and girls at 11 years of age or younger. A Finnish twin study also found that environmental factors shared by co-twins explained a share of BMI variation in children for whom parents had limited education at 11-12 and 14 years of age, but this variance component was not present at 17 years of age (14). Disentangling shared environmental and additive genetic effects requires considerable statistical power (45), and, thus, it is possible that these divergent results within Europe are just because of sample variation. However, these results suggest that in childhood shared environmental factors may contribute directly to the socio-economic differences in BMI, at least in some societies, whereas in adolescence and adulthood, parental education affects BMI mainly by modifying genetic effects.

Our data have strengths but also limitations. The main strength is the large sample size allowing us to construct and test models in narrow age groups and offering the required statistical power to analyze

the effect of parental education over the lifespan. Further, we were able to analyze these differences between cultural-geographic regions representing different levels of obesogenic environment. A limitation is that our data is heavily biased toward Caucasian populations following the Westernized life-style thus limiting statistical power in East Asia. New data collections in East Asia and in other non-westernized cultural contexts would thus be warranted, especially because we found evidence that there may be substantial differences in the effect of parental education on BMI between more and less obesogenic societies. Also our data come from multiple sources, which may differ in representativeness and other factors potentially affecting the strength of found associations.

Conclusion

We found clear evidence that higher parental education is associated with lower BMI and less genetic and environmental variation of BMI over the lifespan. The critical period is located just after early childhood when BMI variation started to increase. In addition to genetic variance, especially in late childhood, the environment shared by co-twins and siblings can be important. The educational differences are particularly prominent in North America and Australia. Socio-economic position of the family affects BMI partly by modifying genetic factors, and the effects on BMI are long lasting, being present through adulthood, especially in obesogenic societies.

References

(1) NCD Risk Factor Collaboration. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet* 2016;387:1377-1396.

(2) Silventoinen K, Jelenkovic A, Sund R, Hur Y, Yokoyama Y, Honda C, et al. Genetic and environmental effects on body mass index from infancy to the onset of adulthood: an individual-based

pooled analysis of 45 twin cohorts participating in the COllaborative project of Development of Anthropometrical measures in Twins (CODATwins) study. *Am J Clin Nutr* 2016;104:371-379.

(3) Silventoinen K, Jelenkovic A, Sund R, Yokoyama Y, Hur YM, Cozen W, et al. Differences in genetic and environmental variation in adult body mass index by sex, age, time period, and region: an individual-based pooled analysis of 40 twin cohorts. *Am J Clin Nutr* 2017;106:457-466.

(4) Turcot V, Lu Y, Highland HM, Schurmann C, Justice AE, Fine RS, et al. Protein-altering variants associated with body mass index implicate pathways that control energy intake and expenditure in obesity. *Nat Genet* 2018;50:26-41.

(5) Yengo L, Sidorenko J, Kemper KE, Zheng Z, Wood AR, Weedon MN, et al. Meta-analysis of genome-wide association studies for height and body mass index in ~700000 individuals of European ancestry. *Hum Mol Genet* 2018;27:3641-3649.

(6) McLaren L. Socioeconomic status and obesity. *Epidemiol Rev* 2007;29:29-48.

(7) Shrewsbury V, Wardle J. Socioeconomic status and adiposity in childhood: a systematic review of cross-sectional studies 1990-2005. *Obesity* 2008;16:275-284.

(8) Fontaine KR, Robertson HT, Holst C, Desmond R, Stunkard AJ, Sørensen TIA, et al. Is socioeconomic status of the rearing environment causally related to obesity in the offspring. *PLoS One* 2011;6:e27692.

(9) Rokholm B, Silventoinen K, Angquist L, Skytthe A, Kyvik KO, Sørensen TIA. Increased genetic variance of BMI with a higher prevalence of obesity. *PLoS One* 2011;6:e20816.

(10) Rokholm B, Silventoinen K, Tynelius P, Gamborg M, Sørensen TIA, Rasmussen F. Increasing genetic variance of body mass index during the Swedish obesity epidemic *PLoS One* 2011;6:e27135.

(11) Silventoinen K, Huppertz C, van Beijsterveldt CE, Bartels M, Willemsen G, Boomsma DI. The genetic architecture of body mass index from infancy to adulthood modified by parental education. *Obesity* 2016;24:2004-2011.

(12) Johnson W, Kyvik KO, Skytthe A, Deary IJ, Sørensen TIA. Education modifies genetic and environmental influences on BMI. *PLoS One* 2011;6(e16290).

(13) Dinescu D, Horn EE, Duncan G, Turkheimer E. Socioeconomic modifiers of genetic and environmental influences on body mass index in adult twins. *Health Psychol* 2016;35:157-166.

(14) Lajunen HR, Kaprio J, Rose RJ, Pulkkinen L, Silventoinen K. Genetic and environmental influences on BMI from late childhood to adolescence are modified by parental education. *Obesity* 2012;20:583-589.

(15) Silventoinen K, Jelenkovic A, Sund R, Honda C, Aaltonen S, Yokoyama Y, et al. The CODATwins Project: The cohort description of collaborative project of development of

anthropometrical measures in twins to study macro-environmental variation in genetic and environmental effects on anthropometric traits. *Twin Res Hum Genet* 2015;18:348-360.

(16) Fielding RA, Vellas B, Evans WJ, Bhasin S, Morley JE, Newman AB, et al. Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. International working group on sarcopenia. *J Am Med Dir Assoc* 2011;12:249-256.

(17) Silventoinen K, Jelenkovic A, Latvala A, Sund R, Yokoyama Y, Ullemar V, et al. Education in twins and their parents across birth cohorts over 100 years: an individual-level pooled analysis of 42 twin cohorts. *Twin Res Human Genet* 2017;20:395-405.

(18) Mallick S, Li H, Lipson M, Mathieson I, Gymrek M, Racimo F, et al. The Simons Genome Diversity Project: 300 genomes from 142 diverse populations. *Nature* 2016;538:201-206.

(19) Posthuma D, Beem AL, de Geus EJ, van Baal GC, von Hjelmberg JB, Iachine I, et al. Theory and practice in quantitative genetics. *Twin Res* 2003;6:361-376.

(20) Jelenkovic A, Yokoyama Y, Sund R, Honda C, Bogl LH, Aaltonen S, et al. Zygosity differences in height and body mass index of twins from infancy to old age: A study of the CODATwins project. *Twin Res Hum Genet* 2015;18:557-570.

(21) Purcell S. Variance components models for gene-environment interaction in twin analysis. *Twin Res* 2002;5:554-571.

(22) Neale MC, Hunter MD, Pritikin JN, Zahery M, Brick TR, Kirkpatrick RM, et al. OpenMx 2.0: Extended structural equation and statistical modeling. *Psychometrika* 2016;81:535-549.

(23) Foraita R, Günther F, Gwozdz W, Reisch LA, Russo P, Lauria F, et al. Does the FTO gene interact with the socioeconomic status on the obesity development among young European children? Results from the IDEFICS study. *Int J Obes* 2015;39:1-6.

(24) Corella D, Carrasco P, Sorli JV, Coltell O, Ortega-Azorín C, Guillén M, et al. Education modulates the association of the FTO rs9939609 polymorphism with body mass index and obesity risk in the Mediterranean population. *Nutr Metab Cardiovasc Dis* 2012;22:651-658.

(25) Pigeyre M, Bokor S, Romon M, Gottrand F, Gilbert CC, Valtueña J, et al. Influence of maternal educational level on the association between the rs3809508 neuromedin B gene polymorphism and the risk of obesity in the HELENA study. *Int J Obes* 2010;34:478-486.

(26) Lagou V, Liu G, Zhu H, Stallmann-Jorgensen IS, Gutin B, Dong Y, et al. Lifestyle and socioeconomic-status modify the effects of ADRB2 and NOS3 on adiposity in European-American and African-American adolescents. *Obesity* 2011;19:595-603.

(27) Bukodi E, Erikson R, Goldthorpe JH. The effects of social origin and cognitive ability on educational attainment: evidence from Britain and Sweden. *Acta Sociologica* 2014;57:293-310.

- (28) Krapohl E, Rimfeld K, Shakeshaft NG, Trzaskowski M, McMillan A, Pingault JB, et al. The high heritability of educational achievement reflects many genetically influenced traits, not just intelligence. *Proc Natl Acad Sci USA* 2014;111:15273-15278.
- (29) Morgen CS, Andersen PK, Mortensen LH, Howe LD, Rasmussen M, Due P, et al. Socioeconomic disparities in birth weight and body mass index during infancy through age 7 years: a study within the Danish National Birth Cohort. *BMJ Open* 2017;7:e011781.
- (30) Howe LD, Tilling K, Galobardes B, Smith GD, Ness AR, Lawlo DA. Socioeconomic disparities in trajectories of adiposity across childhood. *Int J Pediatr Obes* 2011;6:e144-153.
- (31) Bulik-Sullivan B, Finucane HK, Anttila V, Gusev A, Day FR, Loh PR, et al. An atlas of genetic correlations across human diseases and traits. *Nat Genet* 2015;47:1236-1241.
- (32) Sovio U, Mook-Kanamori DO, Warrington NM, Lawrence R, Briollais L, Palmer CN, et al. Association between common variation at the FTO locus and changes in body mass index from infancy to late childhood: the complex nature of genetic association through growth and development. *PLoS Genet* 2011;7(e1001307).
- (33) Elks CE, Loos RJ, Hardy R, Wills AK, Wong A, Wareham NJ, et al. Adult obesity susceptibility variants are associated with greater childhood weight gain and a faster tempo of growth: the 1946 British Birth Cohort Study. *Am J Clin Nutr* 2012;95:1150-1156.
- (34) Warrington NM, Howe LD, Wu YY, Timpson NJ, Tilling K, Pennell CE, et al. Association of a body mass index genetic risk score with growth throughout childhood and adolescence. *PLoS One* 2013;8(e79547).
- (35) Warrington NM, Howe LD, Paternoster L, Kaakinen M, Herrala S, Huikari V, et al. A genome-wide association study of body mass index across early life and childhood. *Int J Epidemiol* 2015;44:700-712.
- (36) Qi Q, Downer MK, Kilpeläinen TO, Taal HR, Barton SJ, Ntalla I. Dietary intake, FTO genetic variants, and adiposity: A combined analysis of over 16,000 children and adolescents. *Diabetes* 2015;64:2467-2476.
- (37) Faith MS, Carnell S, Kral TV. Genetics of food intake self-regulation in childhood: literature review and research opportunities. *Hum Hered* 2014;75:80-89.
- (38) Vainik U, Baker TE, Dadar M, Zeighami Y, Michaud A, Zhang Y, et al. Neurobehavioral correlates of obesity are largely heritable. *Proc Natl Acad Sci USA* 2018;115:9312-9317.
- (39) Gustafson SL, Rhodes RE. Parental correlates of physical activity in children and early adolescents. *Sports Med* 2006;36:79-97.
- (40) Graff M, Scott RA, Justice AE, Young KL, Feitosa MF, Barata L, et al. Genome-wide physical activity interactions in adiposity - A meta-analysis of 200,452 adults. *PLoS Genet* 2017;13(e1006528).

- (41) Silventoinen K, Hasselbalch AL, Lallukka T, Bogl L, Pietiläinen KH, Heitmann BL, et al. Modification effects of physical activity and protein intake on heritability of body size and composition. *Am J Clin Nutr* 2009;90:1096-1103.
- (42) Wilson ME, Fisher J, Fischer A, Lee V, Harris RB, Bartness TJ. Quantifying food intake in socially housed monkeys: social status effects on caloric consumption. *Physiol Behav* 2008;94:586-594.
- (43) Andrews C, Viviani J, Egan E, Bedford T, Brilot B, Nettle D, et al. Early life adversity increases foraging and information gathering in European starlings, *Sturnus vulgaris*. *Anim Behav* 2015;109:123-132.
- (44) Kirk SF, Penney TL, McHugh TL. Characterizing the obesogenic environment: the state of the evidence with directions for future research. *Obesity Reviews* 2009;11:109-117.
- (45) Visscher PM, Gordon S, Neale MC. Power of the classical twin design revisited: II detection of common environmental variance. *Twin Res Hum Genet* 2008;11:48-54.

Figure legends

Figure 1. Mean BMI modification effects of parental education with 95% confidence intervals from 1 until 70-79 years of age by sex.

Figure 2. Mean BMI modification effects of parental education with 95% confidence intervals from 1 until 70-79 years of age by sex and cultural-geographic region.

Figure 3. Total variation of BMI decomposed to additive genetic (gray), shared environmental (white) and unique environmental variations (black) from 1 until 70-79 years of age by sex and parental education.

Figure 4. Total variation of BMI decomposed to additive genetic (gray), shared environmental (white) and unique environmental variations (black) from 1 until 70-79 years of age by parental education and geographic-cultural region in males.

Figure 5. Total variation of BMI decomposed to additive genetic (gray), shared environmental (white) and unique environmental variations (black) from 1 until 70-79 years of age by parental education and geographic-cultural region in females.

Table 1. The number of twin individuals and means and standard deviations (SD) of body mass index (BMI) and parental education by age.¹

Age	N	Males						Females						
		BMI		Maternal education		Paternal education		BMI		Maternal education		Paternal education		
		(kg/m ²)		(years)		(years)		(kg/m ²)		(years)		(years)		
		Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
1	13548	17.1	1.36	13.8	2.31	13.9	2.6	14023	16.7	1.37	13.8	2.33	13.9	2.63
2	11273	16.5	1.37	14.0	2.42	14.0	2.70	11324	16.2	1.37	14.0	2.44	13.9	2.76
3	10907	15.9	1.32	14.2	2.49	14.2	2.73	11460	15.6	1.37	14.2	2.51	14.1	2.76
4	3533	15.9	1.68	15.3	3.39	14.7	3.61	3579	15.7	1.68	15.4	3.38	14.7	3.68
5	6533	15.3	1.58	14.4	2.61	14.3	2.82	6577	15.1	1.66	14.3	2.53	14.3	2.84
6	1898	15.7	1.95	15.1	3.07	14.9	3.23	1998	15.5	1.98	15.2	3.11	14.9	3.32
7	7084	15.4	1.78	14.3	2.44	14.3	2.64	7499	15.4	1.94	14.2	2.48	14.3	2.71
8	4388	15.8	2.06	14.4	2.80	14.3	2.89	4503	15.8	2.22	14.4	2.79	14.3	2.95
9	3497	16.7	2.60	14.7	3.14	14.4	3.23	3471	16.7	2.76	14.8	3.12	14.5	3.28
10	7017	16.6	2.39	14.1	2.57	14.2	2.71	7417	16.7	2.50	14.0	2.46	14.2	2.65
11	3987	17.5	2.69	13.4	3.67	12.9	4.00	4078	17.5	2.92	13.4	3.67	12.9	4.04
12	6760	17.6	2.69	13.8	2.78	13.8	3.13	7023	17.7	2.73	13.7	2.79	13.8	3.18
13	3111	18.6	3.04	14.3	2.87	14.2	3.07	3436	18.7	3.03	14.2	2.81	14.2	3.00
14	5447	19.3	2.94	13.3	3.59	12.9	3.95	6121	19.5	3.02	13.3	3.48	12.9	3.90
15	3278	20.0	3.28	14.3	2.85	14.3	3.02	3603	20.1	3.16	14.2	2.68	14.2	2.93
16	3948	20.8	3.08	13.2	3.19	13.3	3.32	4616	20.6	2.96	13.2	3.07	13.2	3.24
17	5468	21.3	2.91	13.0	3.36	12.8	3.59	6140	20.9	2.95	13.2	3.14	13.0	3.44
18	4224	21.8	2.98	12.0	3.60	11.7	4.10	3775	21.2	3.12	12.7	3.35	12.5	3.74
19	2664	22.0	2.88	12.3	3.04	12.4	3.46	3198	21.5	3.28	13.0	2.78	13.3	3.03
20-29	10005	23.3	3.26	12.0	3.53	11.9	3.91	12096	22.2	3.77	12.5	3.30	12.6	3.62
30-39	7952	25.0	3.45	11.8	2.93	12.0	3.37	11204	23.1	4.18	11.9	2.99	12.2	3.35
40-49	9440	25.7	3.40	11.2	3.22	11.0	3.71	8399	24.3	4.48	11.7	3.00	11.7	3.38
50-59	4720	26.7	3.91	11.0	3.11	10.6	3.54	3274	24.8	4.40	10.8	3.30	10.8	3.41
60-69	4397	26.2	3.54	10.4	3.24	10.0	3.63	2315	24.5	3.90	10.3	3.17	10.3	3.57
70-79	2187	25.6	3.17	10.4	3.40	9.8	3.88	806	24.2	3.93	10.0	3.26	10.4	3.79

¹Data from the following twin projects have been pooled: Europe (Child and Adolescent Twin Study in Sweden, Netherlands Twin Cohort of Children, FinnTwin12, FinnTwin16, East Flanders Prospective Twin Survey, Gemini, Italian Twin Registry, Norwegian Twin Registry, Portugal Twin Cohort, TCHAD-study, Turkish Twin Study); USA (Boston University Twin Project, California Twin Program, Carolina African American Twin Study of Aging, Colorado Twin Registry, Michigan Twins Study, Mid Atlantic Twin Registry, Minnesota Twin Registry, University of Southern California Twin Study, Texas Twin Project, Vietnam Era Twin Study of Aging, NAS-NRC Study); East Asia (Korean Twin-Family Register, Ochanomizu University Twin Project, Qingdao Twin Registry Children, South Korea Twin Registry, West Japan Twins and Higher Order Multiple Births Registry); and others (Australian Twin Registry, Longitudinal Israeli Study of Twins)




Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.