



Impairment of Serotonergic Transmission by the Antiparkinsonian Drug L-DOPA: Mechanisms and Clinical Implications

Cristina Miguelez¹, Abdelhamid Benazzouz^{2,3}, Luisa Ugedo¹ and Philippe De Deurwaerdère^{2,3*}

¹Department of Pharmacology, Faculty of Medicine and Dentistry, University of the Basque Country (UPV/EHU), Leioa, Spain, ²Institut des Maladies Neurodégénératives, Université de Bordeaux, UMR 5293, Bordeaux, France, ³CNRS, Institut des Maladies Neurodégénératives, UMR 5293, Bordeaux, France

The link between the anti-Parkinsonian drug L-3,4-dihydroxyphenylalanine (L-DOPA) and the serotonergic (5-HT) system has been long established and has received increased attention during the last decade. Most studies have focused on the fact that L-DOPA can be transformed into dopamine (DA) and released from 5-HT terminals, which is especially important for the management of L-DOPA-induced dyskinesia. In patients, treatment using L-DOPA also impacts 5-HT neurotransmission; however, few studies have investigated the mechanisms of this effect. The purpose of this review is to summarize the electrophysiological and neurochemical data concerning the effects of L-DOPA on 5-HT cell function. This review will argue that L-DOPA disrupts the link between the electrical activity of 5-HT neurons and 5-HT release as well as that between 5-HT release and extracellular 5-HT levels. These effects are caused by the actions of L-DOPA and DA in 5-HT neurons, which affect 5-HT neurotransmission from the biosynthesis of 5-HT to the impairment of the 5-HT transporter. The interaction between L-DOPA and 5-HT transmission is especially relevant in those Parkinson's disease (PD) patients that suffer dyskinesia, comorbid anxiety or depression, since the efficacy of antidepressants or 5-HT compounds may be affected.

Keywords: serotonin, dopamine, electrophysiology, intracerebral microdialysis, depression, dyskinesia, Parkinson's disease, exocytosis

INTRODUCTION

Parkinson's disease (PD) is a neurodegenerative disorder characterized by motor symptoms such as bradykinesia, rigidity, resting tremor, postural abnormalities and gait deficits. However, most patients also develop additional non-motor symptoms, such as anxiety, depression, fatigue, apathy, mild cognitive disturbances and dementia (Bastide et al., 2015). The disease is no longer seen as a specific consequence of dopaminergic (DA) neurodegeneration, since other neuronal systems, such as the noradrenergic and serotonergic (5-HT) systems, also suffer alterations in the course of the disease (Jenner et al., 1983; Delaville et al., 2011). The 5-HT system is involved

Abbreviations: 5-HT, serotonin; 5-HTTP, 5-hydroxytryptophan; 5-HIAAL, 5-hydroxyindole acetaldehyde; AADC, amino acid decarboxylase; DA, dopamine; DRN, dorsal raphe nucleus; HIPP, hippocampus; L-DOPA, L-3,4-dihydroxyphenylalanine; MAO, monoamine oxidase; PD, Parkinson's disease; PFC, prefrontal cortex; SERT, serotonin transporter; SNr, substantia nigra reticulata; SSRI, selective serotonin reuptake inhibitors; STR, striatum; VMAT2, monoamine vesicular transporter.

OPEN ACCESS

Edited by:

Yu-Qiang Ding,
Tongji University, China

Reviewed by:

Carla Cannizzaro,
University of Palermo, Italy
Yue Wang,
Shandong University, China

*Correspondence:

Phillippe De Deurwaerdère
deurwaer@u-bordeaux2.fr

Received: 30 March 2017

Accepted: 25 August 2017

Published: 12 September 2017

Citation:

Miguelez C, Benazzouz A, Ugedo L and De Deurwaerdère P (2017) Impairment of Serotonergic Transmission by the Antiparkinsonian Drug L-DOPA: Mechanisms and Clinical Implications. *Front. Cell. Neurosci.* 11:274. doi: 10.3389/fncel.2017.00274

in some PD pathological clinical manifestations as well as in some side effects induced by long-term treatment using L-3,4-dihydroxyphenylalanine (L-DOPA), such as L-DOPA-induced dyskinesia (Scholtissen et al., 2006a,b; Haduch et al., 2016). It has been suggested that 5-HT neurons are mainly responsible for the increase in DA release induced by chronic L-DOPA treatment in the striatum (Carta et al., 2007, 2008). However, the motor improvements or side effects produced by low doses of L-DOPA (3–6 mg/kg) cannot be explained only by what happens in the striatum since those doses fail to produce physiological levels of DA in that nucleus (Navailles et al., 2010b, 2011; Nevalainen et al., 2011; Porras et al., 2014; De Deurwaerdère et al., 2017). Indeed, L-DOPA may act through other mechanisms (Misu et al., 1996; Porras et al., 2014; De Deurwaerdère et al., 2017) or modify DA transmission in other brain regions to a greater extent than in the striatum (Navailles and De Deurwaerdère, 2012b). However, despite the evidence that L-DOPA alters the 5-HT system (Navailles and De Deurwaerdère, 2012a), and that 5-HT receptor agonists can produce therapeutic benefit in L-DOPA-induced dyskinesia and psychosis, the impact of L-DOPA on 5-HT neurotransmission has been poorly studied (De Deurwaerdère et al., 2017).

The purpose of this mini-review is to assess the experimental data highlighting the complex effects of L-DOPA on 5-HT extracellular levels and ultimately on 5-HT transmission.

SEROTONERGIC NEURONS MEDIATE L-DOPA-DERIVED DA RELEASE

L-DOPA can be taken up by virtually all cells in the brain and decarboxylated into DA in cells expressing amino acid decarboxylase (AADC) and other types of decarboxylase such as histidine decarboxylase (De Deurwaerdère et al., 2017). The reason why 5-HT neurons differ from other cell types expressing AADC is that in 5-HT neurons newly synthesized DA can directly compete with 5-HT for inclusion in exocytotic vesicles via the vesicular monoamine transporter VMAT2 (Lohoff, 2010; De Deurwaerdère et al., 2017). L-DOPA-induced DA is thereby concentrated in 5-HT vesicles, probably at the expense of endogenous 5-HT (see below), and is released by 5-HT neurons in well-innervated areas of the whole brain (Navailles et al., 2010b). However, 5-HT cells lack the mechanisms to control DA release, as 5-HT_{1A} and 5-HT_{1B} autoreceptors and 5-HT transporters (SERT) do not detect extracellular DA. Several microdialysis studies show that L-DOPA, at 12 mg/kg and higher doses, induces excessive DA release (Abercrombie et al., 1990; Kannari et al., 2000; Navailles et al., 2010b). However, the amount of released DA may have been overestimated in animal models of PD, as there are no clearance mechanisms and consequently extracellular L-DOPA-derived DA is detected by probes in significant quantities (Miller and Abercrombie, 1999). In fact, the real levels of extracellular DA in the striatum are probably much lower than those measured, given the low density of 5-HT terminals in the striatum (De Deurwaerdère et al., 2017).

In general, 5-HT neurons have the required enzymatic/transporter equipment to transform L-DOPA

into DA and release it. However, 5-HT neurons lack the proper autoregulatory mechanisms to control DA release and clearance, which can cause excessive DA levels in several brain regions.

EFFECT OF L-DOPA ON THE ELECTRICAL ACTIVITY OF 5-HT NEURONS

Several studies suggest that L-DOPA could modify the activity of 5-HT neurons. First, DA agonists increase dorsal raphe nucleus (DRN) activity in control animals (Martín-Ruiz et al., 2001); second, DA neurons degeneration unevenly alters the electrophysiological characteristic of DRN neurons (Zhang et al., 2007; Guiard et al., 2008; Kaya et al., 2008; Wang et al., 2009; Prinz et al., 2013) and third, DA and 5-HT release from 5-HT cell bodies and terminals may also impact 5-HT system activity. A computational study has modeled the latter impact, which predicts that L-DOPA administration will decrease 5-HT release in the DRN and subsequently increase DRN neuron firing rate (Reed et al., 2012).


Patch clamp recordings performed in brain slices from control animals with intact DA innervation demonstrated that acute L-DOPA depresses 5-HT_{1A} receptor-mediated transmission in the DRN (Gantz et al., 2015) and discretely increases the basal firing activity of 5-HT neurons (Prinz et al., 2013). On the other hand, in a study performed in anesthetized rats, systemic administration of therapeutic doses of L-DOPA did not modify DRN activity (Migueluez et al., 2013).

In animal models of PD, neither acute nor chronic systemic administration of L-DOPA altered the neuronal activity of DRN cells (Migueluez et al., 2016a,b). *In vitro*, L-DOPA partially reversed the changes in excitability observed in Parkinsonian mice (Prinz et al., 2013).

In conclusion, the effect of L-DOPA on DRN neuron activity seems non-existent or discrete, suggesting that the interaction between the 5-HT system and L-DOPA may be more relevant at the terminal rather than the somatic level.

EFFECT OF L-DOPA ON 5-HT EXTRACELLULAR LEVELS

As discussed above, L-DOPA-derived DA competes with 5-HT to enter exocytotic vesicles at 5-HT terminals, which likely leads to an inhibition of 5-HT release. However, this hypothesis has not been fully supported by *in vitro* and *in vivo* data, which suggests that the effects of L-DOPA on extracellular 5-HT levels are more complex than previously thought. Indeed, when applied locally, L-DOPA enhances the 5-HT efflux (Biggs and Starr, 1999) while when administered systemically, it either does not alter or reduces 5-HT extracellular levels (Lindgren et al., 2010; Navailles et al., 2010a, 2014; Navailles and De Deurwaerdère, 2012a). Furthermore, L-DOPA's effects on extracellular 5-HT levels are not similar in all regions innervated by the DRN including no effect (striatum), inhibition (substantia nigra and cortex) and biphasic excitation/inhibition (hippocampus;



Navailles et al., 2011; Navailles and De Deurwaerdère, 2012b; **Figure 1**).


In accordance with the idea that newly synthesized DA displaces 5-HT inside exocytotic vesicles, several publications show that acute administration of L-DOPA reduces 5-HT tissue content, which mainly represents 5-HT stored in vesicle compartments, or its metabolite 5-hydroxyindolacetic acid (Eskow Jaunarajs et al., 2012; Migueluez et al., 2016b). Nonetheless, some discrepancies exist which could depend on the doses used of L-DOPA or AADC inhibitor, or the time of sacrifice after L-DOPA administration. After chronic administration of L-DOPA (12 mg/kg), L-DOPA further inhibited extracellular 5-HT levels in the hippocampus (amplified the inhibitory component of the biphasic effect) and substantia nigra, while its effect was unchanged in the cortex and the striatum (Navailles et al., 2011). In control and Parkinsonian monkeys, the acute administration of L-DOPA reduces 5-HT tissue levels in the striatum and motor cortex. In contrast, chronic L-DOPA treatment reduced 5-HT levels in the striatum, hippocampus or amygdala of Parkinsonian but not control monkeys (Engeln et al., 2015). In Parkinsonian rats, chronic treatment using 6 mg/kg of L-DOPA reduces 5-HT content in a region-dependent manner (Stansley and Yamamoto, 2014, 2015a).

In summary, although local administration of L-DOPA may increase 5-HT levels, systemic administration reduces it in several brain regions without modifying DRN neuron activity and modestly reducing 5-HT extracellular levels.

THE ROLE OF NON-EXOCYTOTIC MECHANISMS IN THE EFFECT OF L-DOPA

As anticipated as early as 1970 (Ng et al., 1970), apart from exocytotic release, L-DOPA also triggers non-exocytotic (non-vesicular) efflux of 5-HT and DA. Indeed, supra-therapeutic dose [100 mg/kg (De Deurwaerdère et al., 2017)] of L-DOPA transiently enhances 5-HT release. A Ca^{2+} free medium magnified this effect and unmasked excitation of 5-HT release induced by a therapeutic dose (12 mg/kg) of L-DOPA (Migueluez et al., 2016b). In addition, blockade of action potential-dependent presynaptic release or Ca^{2+} removal in the perfusion solution does not suppress L-DOPA-induced DA release (Miller and Abercrombie, 1999; Lindgren et al., 2010; Migueluez et al., 2016b).

The mechanism underlying the non-exocytotic release of neurotransmitters is still unclear. It may also require 5-HT neurons (Tanaka et al., 1999; Navailles et al., 2010b) and the involvement of one or several transporters, but the effect also



shows region dependency. Indeed, SERT blockade by citalopram significantly reduced peripheral L-DOPA-induced DA release in the hippocampus but not in the prefrontal cortex (PFC; Migueluez et al., 2016b; **Figure 2**).

The participation of a non-vesicular release of both DA and 5-HT in different brain regions may contribute to the heterogeneous effect of L-DOPA on these neurotransmitters. The whole biochemical picture appears to be particularly complicated for 5-HT as it is displaced by DA in exocytotic vesicles, trapped in the neuron by its reduced ability to exit via the SERT, and is perhaps more available to degradation by monoamine oxidase (**Figure 2**; Stansley and Yamamoto, 2014). Moreover, the metabolism of both DA and 5-HT could have detrimental effects on the survival of groups of 5-HT neurons (Navailles et al., 2011; Stansley and Yamamoto, 2015a). Indeed, L-DOPA can promote oxidative stress and generate free radicals or toxic compounds. Therefore, L-DOPA-derived DA can be metabolized into pro-oxidant quinones, 3,4-dihydroxyphenylacetic acid or hydrogen peroxide and produce cell death in the DRN (Stansley and Yamamoto, 2015b).

To sum up, *in vitro* experiments and mathematical models predict L-DOPA function in 5-HT terminals (Reed et al., 2012) but none of these models fully describe the region dependency and the complex modalities of release induced by L-DOPA *in vivo*.

ALTERED REACTIVITY OF 5-HT NEURONS IN THE PRESENCE OF L-DOPA

After chronic treatment using L-DOPA and/or DA denervation, the inhibitory control exerted by 5-HT_{1A} receptors over the electrical activity of 5-HT neurons is maintained (Migueluez et al., 2016a). In these conditions, 5-HT_{1A} receptor agonists completely inhibit DRN neuron activity and diminish L-DOPA-induced DA release at the terminal sites (Kannari et al., 2001; Iderberg et al., 2015). However, while the firing rate is completely suppressed, DA release still occurs. This result is compatible with the evidence that some mechanisms related to DA release are independent of the electrical activity of 5-HT neurons. The control exerted by 5-HT_{1B} receptors at the terminals might not be fully functional by itself because the stimulation of these receptors alone does not reduce L-DOPA-stimulated striatal DA release (Kannari et al., 2001) but ameliorates dyskinesia when co-administered together with 5-HT_{1A} agonists (Muñoz et al., 2009). The activation of other 5-HT receptors can also modulate L-DOPA-stimulated DA release. In this regard, 5-HT₄ receptor stimulation can indirectly activate the activity of 5-HT neurons (Ge and Barnes, 1996; Lucas and Debonnel, 2002; Lucas et al., 2005, 2007) and enhance L-DOPA-induced DA release in the PFC and the substantia nigra, but not in the hippocampus or the striatum (Navailles et al., 2015).

In Parkinsonian conditions, SERT density might not be altered but L-DOPA-derived DA can directly and indirectly modify SERT function. As reported above, DA can also bind to SERT and compete with 5-HT for non-exocytotic efflux through this transporter (**Figure 2**). The indirect impairment is related to a possible decrease in 5-HT cellular levels. Indeed, fluoxetine or

citalopram can reduce L-DOPA-stimulated DA release (Yamato et al., 2001; Navailles et al., 2010b) through a mechanism partly involving 5-HT_{1A} receptors (Yamato et al., 2001). However, the efficacy of fluoxetine to inhibit the electrical activity of 5-HT neurons in the DRN can be reduced by L-DOPA, in line with a possible inhibitory effect of L-DOPA on extracellular levels of 5-HT (Migueluez et al., 2016a). Behavioral studies support a lower efficacy of selective serotonin reuptake inhibitors (SSRIs) in the presence of L-DOPA (Migueluez et al., 2013; Fidalgo et al., 2015). Because the SERT plays a critical role in the various outcomes of L-DOPA-induced changes of DA and 5-HT extracellular levels, its indirect modulation by 5-HT_{2B} receptor ligands can be tested. Indeed, 5-HT_{2B} receptors can be considered as autoreceptors involved in the control of SERT activity and their antagonism has been shown to block outward release induced by the 5-HT/DA releaser 3,4-methylenedioxymethamphetamine (Gudelsky and Yamamoto, 2008).

Extracellular levels of 5-HT in the presence of L-DOPA can be impacted in different ways by: (1) decreased exocytosis, which non-exocytotic release can compensate for to a limited extent; (2) the loss of extracellular clearance by the SERT (competition with DA); and (3) the loss of other clearance mechanisms such as the noradrenaline transporter (Navailles et al., 2014). On the basis of the described data, it seems that extracellular levels of 5-HT are almost impossible to control in a context where all these factors are region dependent. For instance, the inhibition of levels of extracellular 5-HT induced by the high frequency stimulation of the subthalamic nucleus was attenuated by the presence of L-DOPA (technically an increase compared with what was expected), while L-DOPA-stimulated DA extracellular levels were partly decreased (Navailles et al., 2010a). L-DOPA disrupts the control of extracellular 5-HT levels, and the biological function of this phenomenon remains completely unknown.

In summary, the release of L-DOPA-derived DA from 5-HT neurons is a complicated mechanism involving 5-HT receptors and transporters in a region-dependent manner.

FROM THE NEUROCHEMISTRY TO THE CLINIC

The side effects induced by L-DOPA are usually interpreted as a consequence of the modification of L-DOPA-stimulated DA release from 5-HT neurons involving alteration of 5-HT neurotransmission. Although the published results are heterogeneous, it is accepted that L-DOPA induces adaptive and/or toxic changes in the 5-HT system that may have clinical relevance. One study in postmortem tissue described a positive correlation between SERT binding densities and the development of dyskinesia (Rylander et al., 2010); however, subsequent studies in postmortem samples or patients did not find any correlation between 5-HT or SERT in the striatum and dyskinesia (Politis et al., 2014; Cheshire et al., 2015). These discrepant results may be related to the overestimated role of the striatum in dyskinesia (De Deurwaerdère et al., 2017). Indeed, several brain regions other than the striatum have been proposed to participate in dyskinesia. A recent publication showed that SERT binding was positively correlated with the manifestation

of dyskinesia in the internal and external parts of the globus pallidus (Smith et al., 2015). Regarding 5-HT receptors, PD is marked by the uneven modification of some 5-HT receptor subtypes in different brain regions. Therefore, Parkinsonian patients treated using L-DOPA show higher expression of 5-HT_{1A} and 5-HT_{2C} receptors in the cerebral cortex, but not in other brain regions (Huot et al., 2010, 2012).

The clinical implications of these 5-HT adaptive changes could underlie the efficacy of drugs acting on 5-HT neurotransmission to reduce the severity of dyskinesia. Although in animal models of the disease several drugs acting on 5-HT_{1A/1C} or 5-HT_{1B} receptors decrease the expression of L-DOPA-induced dyskinesia (Migueluez et al., 2013), translational extrapolation to the clinic has been less positive. So far, only two drugs have been proven to be clinically efficient in treating dyskinesia. These are eltoprazine, a 5-HT_{1A/B} receptor agonist (Bomasang-Layno et al., 2015; Svenningsson et al., 2015), and buspirone, a partial agonist of the 5-HT_{1A} receptor, which discretely ameliorates dyskinesia (Politis et al., 2014). In general, 5-HT compounds have been demonstrated to be less efficacious than expected, probably because the existing theories about 5-HT and DA interaction, especially in the striatum, are too simplistic.

Another important clinical aspect to take into account is the fact that often antidepressants and L-DOPA are administered together. Commonly used SSRIs enhance 5-HT transmission in depressed patients; however, in PD the DRN undergoes degeneration contributing to worse therapeutic control of depressive symptoms in these patients (Deurwaerdère and Ding, 2016). Preclinical data show a loss of efficacy of SSRIs and suggest the use of other strategies such as noradrenaline uptake inhibitors (Migueluez et al., 2013). Unfortunately, the clinical data are unclear. Some studies suggest that SSRIs are less efficacious for treating depression in PD and should be used only as a last choice (Aarsland et al., 2009; Skapinakis et al., 2010; Liu et al., 2013; Rocha et al., 2013). However, a recent meta-analysis reports that treatment using SSRIs significantly improves depression among PD patients (Bomasang-Layno et al., 2015). Other antidepressants, such as the mixed noradrenaline and 5-HT uptake inhibitor, venlafaxine, had similar efficacy as the SSRI paroxetine (Richard et al., 2012; Broen et al., 2016). However, more studies should be conducted to verify which antidepressant has a better therapeutic profile when specifically co-administered with L-DOPA, which is rarely evaluated.

L-DOPA can also produce psychosis in the advanced stages of the PD, although less frequency than dyskinesia. The

manifestation of psychosis could depend on an excess of DA transmission, presumably in cortical areas (De Deurwaerdère and Di Giovanni, 2017), and drugs that limit the excess of DA release from 5-HT neurons could be therapeutically interesting. Therefore, the atypical antipsychotic drugs clozapine, risperidone and olanzapine that interact with 5-HT_{1A}, 5-HT_{2A} and 5-HT_{2C} receptors are efficacious even at doses lower than those classically used in the treatment of schizophrenia. The 5-HT_{2A} inverse agonist primavanserin, recently approved to treat PD psychosis, has shown discrete therapeutical improvements in clinical trials although more postmarketing studies are necessary (Meltzer et al., 2010; Cummings et al., 2014; Divac et al., 2016). In terms of mechanism of action, it is unclear whether primavanserin reduces 5-HT tone at 5-HT_{2A} receptors and/or stabilizes DA transmission.

CONCLUSION

The effects of L-DOPA on both DA and 5-HT extracellular levels involve 5-HT neurons. These effects are complex and involve several mechanisms. One of the most surprising outcomes is the region-dependent effect for both neurotransmitters. Regarding 5-HT levels, there is a disparity between the results from electrophysiological, biochemical and neurochemical studies. L-DOPA affects 5-HT neurotransmission in the brain through the numerous actions of L-DOPA and DA inside 5-HT neurons, i.e., effects on 5-HT biosynthesis or on 5-HT transporter level and neuron survival. Importantly, the impact of L-DOPA on 5-HT transmission concerns the therapeutic effects of 5-HT drugs on L-DOPA-induced dyskinesia, comorbid anxiety and depression in PD patients.

AUTHOR CONTRIBUTIONS

CM and PDD have proposed the architecture of the article and proposed the draft that was implemented by LU and AB. CM and PDD designed the figure and edited the manuscript.

FUNDING

The project was funded by grants from the Government of the Basque Country (IT 747-13), the Spanish Government [SAF2016-77758-R (AEI/FEDER, UE)] and by the Fondation de France. PDD acknowledges the support given by the cooperation for science and technology (COST) action CM15120. PDD and AB acknowledge the support of the Centre National de la Recherche Scientifique.

REFERENCES

- Aarsland, D., Marsh, L., and Schrag, A. (2009). Neuropsychiatric symptoms in Parkinson's disease. *Mov. Disord.* 24, 2175–2186. doi: 10.1002/mds.22589
- Abercrombie, E. D., Bonatz, A. E., and Zigmond, M. J. (1990). Effects of L-dopa on extracellular dopamine in striatum of normal and 6-hydroxydopamine-treated rats. *Brain Res.* 525, 36–44. doi: 10.1016/0006-8993(90)91318-b
- Bastide, M. F., Meissner, W. G., Picconi, B., Fasano, S., Fernagut, P.-O., Feyder, M., et al. (2015). Pathophysiology of L-dopa-induced motor and non-motor complications in Parkinson's disease. *Prog. Neurobiol.* 132, 96–168. doi: 10.1016/j.pneurobio.2015.07.002
- Biggs, C. S., and Starr, M. S. (1999). Microdialysis study of the effects of the antiparkinsonian drug bupropion on L-DOPA-induced release of dopamine and 5-hydroxytryptamine by rat substantia nigra and corpus striatum. *Synapse* 34, 36–46. doi: 10.1002/(sici)1098-2396(199910)34:1<36::aid-syn5>3.3.co;2-7
- Bomasang-Layno, E., Fadlon, I., Murray, A. N., and Himelhoch, S. (2015). Antidepressive treatments for Parkinson's disease: a systematic review and

- meta-analysis. *Parkinsonism Relat. Disord.* 21, 833–842; discussion 833. doi: 10.1016/j.parkreldis.2015.04.018
- Broen, M. P. G., Leentjens, A. F. G., Köhler, S., Kuijff, M. L., McDonald, W. M., and Richard, I. H. (2016). Trajectories of recovery in depressed Parkinson's disease patients treated with pramipexole or venlafaxine. *Parkinsonism Relat. Disord.* 23, 80–85. doi: 10.1016/j.parkreldis.2015.12.013
- Carta, M., Carlsson, T., Kirik, D., and Björklund, A. (2007). Dopamine released from 5-HT terminals is the cause of L-DOPA-induced dyskinesia in parkinsonian rats. *Brain* 130, 1819–1833. doi: 10.1093/brain/awm082
- Carta, M., Carlsson, T., Muñoz, A., Kirik, D., and Björklund, A. (2008). Serotonin-dopamine interaction in the induction and maintenance of L-DOPA-induced dyskinesias. *Prog. Brain Res.* 172, 465–478. doi: 10.1016/s0079-6123(08)00922-9
- Cheshire, P., Ayton, S., Bertram, K. L., Ling, H., Li, A., McLean, C., et al. (2015). Serotonergic markers in Parkinson's disease and levodopa-induced dyskinesias. *Mov. Disord.* 30, 796–804. doi: 10.1002/mds.26144
- Cummings, J., Isaacson, S., Mills, R., Williams, H., Chi-Burris, K., Corbett, A., et al. (2014). Pimavanserin for patients with Parkinson's disease psychosis: a randomised, placebo-controlled phase 3 trial. *Lancet* 383, 533–540. doi: 10.1016/S0140-6736(13)62106-6
- De Deurwaerdère, P., and Di Giovanni, G. (2017). Serotonergic modulation of the activity of mesencephalic dopaminergic systems: therapeutic implications. *Prog. Neurobiol.* 151, 175–236. doi: 10.1016/j.pneurobio.2016.03.004
- De Deurwaerdère, P., Di Giovanni, G., and Millan, M. J. (2017). Expanding the repertoire of L-DOPA's actions: a comprehensive review of its functional neurochemistry. *Prog. Neurobiol.* 151, 57–100. doi: 10.1016/j.pneurobio.2016.07.002
- Delaville, C., Deurwaerdère, P. D., and Benazzouz, A. (2011). Noradrenaline and Parkinson's disease. *Front. Syst. Neurosci.* 5:31. doi: 10.3389/fnsys.2011.00031
- Deurwaerdère, P. D., and Ding, Y. (2016). Antiparkinsonian treatment for depression in Parkinson's disease: are selective serotonin reuptake inhibitors recommended? *Trans. Neurosci. Clin.* 2, 138–149. doi: 10.18679/CN11-6030/R.2016.019
- Divac, N., Stojanović, R., Savić Vujović, K., Medić, B., Damjanović, A., and Prostran, M. (2016). The efficacy and safety of antipsychotic medications in the treatment of psychosis in patients with Parkinson's disease. *Behav. Neurol.* 2016:4938154. doi: 10.1155/2016/4938154
- Engeln, M., De Deurwaerdère, P., Li, Q., Bezaud, E., and Fernagut, P.-O. (2015). Widespread monoaminergic dysregulation of both motor and non-motor circuits in Parkinsonism and dyskinesia. *Cereb. Cortex* 25, 2783–2792. doi: 10.1093/cercor/bhu076
- Eskow Jaunarajs, K. L., George, J. A., and Bishop, C. (2012). L-DOPA-induced dysregulation of extrastriatal dopamine and serotonin and affective symptoms in a bilateral rat model of Parkinson's disease. *Neuroscience* 218, 243–256. doi: 10.1016/j.neuroscience.2012.05.052
- Fidalgo, C., Ko, W. K. D., Tronci, E., Li, Q., Stancampiano, R., Chuan, Q., et al. (2015). Effect of serotonin transporter blockade on L-DOPA-induced dyskinesia in animal models of Parkinson's disease. *Neuroscience* 298, 389–396. doi: 10.1016/j.neuroscience.2015.04.027
- Gantz, S. C., Levitt, E. S., Llamas, N., Neve, K. A., and Williams, J. T. (2015). Depression of serotonin synaptic transmission by the dopamine precursor L-DOPA. *Cell Rep.* 12, 944–954. doi: 10.1016/j.celrep.2015.07.005
- Ge, J., and Barnes, N. M. (1996). 5-HT₄ receptor-mediated modulation of 5-HT release in the rat hippocampus *in vivo*. *Br. J. Pharmacol.* 117, 1475–1480. doi: 10.1111/j.1476-5381.1996.tb15309.x
- Gudelsky, G. A., and Yamamoto, B. K. (2008). Actions of 3,4-methylenedioxymethamphetamine (MDMA) on cerebral dopaminergic, serotonergic and cholinergic neurons. *Pharmacol. Biochem. Behav.* 90, 198–207. doi: 10.1016/j.pbb.2007.10.003
- Guiard, B. P., El Mansari, M., Merali, Z., and Blier, P. (2008). Functional interactions between dopamine, serotonin and norepinephrine neurons: an *in-vivo* electrophysiological study in rats with monoaminergic lesions. *Int. J. Neuropsychopharmacol.* 11, 625–639. doi: 10.1017/s1461145707008383
- Haduch, A., Bromek, E., Wójcikowski, J., Golembiowska, K., and Daniel, W. A. (2016). Melatonin supports CYP2D-mediated serotonin synthesis in the brain. *Drug Metab. Dispos.* 44, 445–452. doi: 10.1124/dmd.115.067413
- Huot, P., Johnston, T. H., Darr, T., Hazrati, L.-N., Visanji, N. P., Pires, D., et al. (2010). Increased 5-HT_{2A} receptors in the temporal cortex of parkinsonian patients with visual hallucinations. *Mov. Disord.* 25, 1399–1408. doi: 10.1002/mds.23083
- Huot, P., Johnston, T. H., Winkelmolen, L., Fox, S. H., and Brotchie, J. M. (2012). 5-HT_{2A} receptor levels increase in MPTP-lesioned macaques treated chronically with L-DOPA. *Neurobiol. Aging* 33, 194.e5–194.e15. doi: 10.1016/j.neurobiolaging.2010.04.035
- Iderberg, H., McCreary, A. C., Varney, M. A., Kleven, M. S., Koek, W., Bardin, L., et al. (2015). NLX-112, a novel 5-HT_{1A} receptor agonist for the treatment of L-DOPA-induced dyskinesia: behavioral and neurochemical profile in rat. *Exp. Neurol.* 271, 335–350. doi: 10.1016/j.expneurol.2015.05.021
- Jenner, P., Sheehy, M., and Marsden, C. D. (1983). Noradrenaline and 5-hydroxytryptamine modulation of brain dopamine function: implications for the treatment of Parkinson's disease. *Br. J. Clin. Pharmacol.* 15, 277S–289S. doi: 10.1111/j.1365-2125.1983.tb05876.x
- Kannari, K., Tanaka, H., Maeda, T., Tomiyama, M., Suda, T., and Matsunaga, M. (2000). Reserpine pretreatment prevents increases in extracellular striatal dopamine following L-DOPA administration in rats with nigrostriatal denervation. *J. Neurochem.* 74, 263–269. doi: 10.1046/j.1471-4159.2000.0740263.x
- Kannari, K., Yamato, H., Shen, H., Tomiyama, M., Suda, T., and Matsunaga, M. (2001). Activation of 5-HT_{1A} but not 5-HT_{1B} receptors attenuates an increase in extracellular dopamine derived from exogenously administered L-DOPA in the striatum with nigrostriatal denervation. *J. Neurochem.* 76, 1346–1353. doi: 10.1046/j.1471-4159.2001.00184.x
- Kaya, A. H., Vlamings, R., Tan, S., Lim, L. W., Magill, P. J., Steinbusch, H. W. M., et al. (2008). Increased electrical and metabolic activity in the dorsal raphe nucleus of Parkinsonian rats. *Brain Res.* 1221, 93–97. doi: 10.1016/j.brainres.2008.05.019
- Lindgren, H. S., Andersson, D. R., Lagerkvist, S., Nissbrandt, H., and Cenci, M. A. (2010). L-DOPA-induced dopamine efflux in the striatum and the substantia nigra in a rat model of Parkinson's disease: temporal and quantitative relationship to the expression of dyskinesia. *J. Neurochem.* 112, 1465–1476. doi: 10.1111/j.1471-4159.2009.06556.x
- Liu, J., Dong, J., Wang, L., Su, Y., Yan, P., and Sun, S. (2013). Comparative efficacy and acceptability of antidepressants in Parkinson's disease: a network meta-analysis. *PLoS One* 8:e76651. doi: 10.1371/journal.pone.0076651
- Lohoff, F. W. (2010). Genetic variants in the vesicular monoamine transporter 1 (VMAT1/SLC18A1) and neuropsychiatric disorders. *Methods Mol. Biol.* 637, 165–180. doi: 10.1007/978-1-60761-700-6_9
- Lucas, G., Compan, V., Charnay, Y., Neve, R. L., Nestler, E. J., Bockaert, J., et al. (2005). Frontocortical 5-HT₄ receptors exert positive feedback on serotonergic activity: viral transfections, subacute and chronic treatments with 5-HT₄ agonists. *Biol. Psychiatry* 57, 918–925. doi: 10.1016/j.biopsych.2004.12.023
- Lucas, G., and Debonnel, G. (2002). 5-HT₄ receptors exert a frequency-related facilitatory control on dorsal raphe nucleus 5-HT neuronal activity. *Eur. J. Neurosci.* 16, 817–822. doi: 10.1046/j.1460-9568.2002.02150.x
- Lucas, G., Rymar, V. V., Du, J., Mnie-Filali, O., Bisgaard, C., Manta, S., et al. (2007). Serotonin(4) (5-HT₄) receptor agonists are putative antidepressants with a rapid onset of action. *Neuron* 55, 712–725. doi: 10.1016/j.neuron.2007.07.041
- Martín-Ruiz, R., Puig, M. V., Celada, P., Shapiro, D. A., Roth, B. L., Mengod, G., et al. (2001). Control of serotonergic function in medial prefrontal cortex by serotonin-2A receptors through a glutamate-dependent mechanism. *J. Neurosci.* 21, 9856–9866.
- Meltzer, H. Y., Mills, R., Revell, S., Williams, H., Johnson, A., Bahr, D., et al. (2010). Pimavanserin, a serotonin(2A) receptor inverse agonist, for the treatment of parkinson's disease psychosis. *Neuropsychopharmacology* 35, 881–892. doi: 10.1038/npp.2009.176
- Migueluez, C., Berrocoso, E., Mico, J. A., and Ugedo, L. (2013). L-DOPA modifies the antidepressant-like effects of reboxetine and fluoxetine in rats. *Neuropharmacology* 67, 349–358. doi: 10.1016/j.neuropharm.2012.11.016
- Migueluez, C., Navailles, S., De Deurwaerdère, P., and Ugedo, L. (2016a). The acute and long-term L-DOPA effects are independent from changes in the activity of dorsal raphe serotonergic neurons in 6-OHDA lesioned rats. *Br. J. Pharmacol.* 173, 2135–2146. doi: 10.1111/bph.13447
- Migueluez, C., Navailles, S., Delaville, C., Marquis, L., Lagière, M., Benazzouz, A., et al. (2016b). L-DOPA elicits non-vesicular releases of serotonin and dopamine in hemiparkinsonian rats *in vivo*. *Eur. Neuropsychopharmacol.* 26, 1297–1309. doi: 10.1016/j.euroneuro.2016.05.004

- Miller, D. W., and Abercrombie, E. D. (1999). Role of high-affinity dopamine uptake and impulse activity in the appearance of extracellular dopamine in striatum after administration of exogenous L-DOPA: studies in intact and 6-hydroxydopamine-treated rats. *J. Neurochem.* 72, 1516–1522. doi: 10.1046/j.1471-4159.1999.721516.x
- Misu, Y., Goshima, Y., Ueda, H., and Okamura, H. (1996). Neurobiology of L-DOPAergic systems. *Prog. Neurobiol.* 49, 415–454. doi: 10.1016/0301-0082(96)00025-1
- Muñoz, A., Carlsson, T., Tronci, E., Kirik, D., Björklund, A., and Carta, M. (2009). Serotonin neuron-dependent and -independent reduction of dyskinesia by 5-HT1A and 5-HT1B receptor agonists in the rat Parkinson model. *Exp. Neurol.* 219, 298–307. doi: 10.1016/j.expneurol.2009.05.033
- Navailles, S., Benazzouz, A., Bioulac, B., Gross, C., and De Deurwaerdère, P. (2010a). High-frequency stimulation of the subthalamic nucleus and L-3,4-dihydroxyphenylalanine inhibit *in vivo* serotonin release in the prefrontal cortex and hippocampus in a rat model of Parkinson's disease. *J. Neurosci.* 30, 2356–2364. doi: 10.1523/jneurosci.5031-09.2010
- Navailles, S., Bioulac, B., Gross, C., and De Deurwaerdère, P. (2010b). Serotonergic neurons mediate ectopic release of dopamine induced by L-DOPA in a rat model of Parkinson's disease. *Neurobiol. Dis.* 38, 136–143. doi: 10.1016/j.nbd.2010.01.012
- Navailles, S., Bioulac, B., Gross, C., and De Deurwaerdère, P. (2011). Chronic L-DOPA therapy alters central serotonergic function and L-DOPA-induced dopamine release in a region-dependent manner in a rat model of Parkinson's disease. *Neurobiol. Dis.* 41, 585–590. doi: 10.1016/j.nbd.2010.11.007
- Navailles, S., and De Deurwaerdère, P. (2012a). Contribution of serotonergic transmission to the motor and cognitive effects of high-frequency stimulation of the subthalamic nucleus or levodopa in Parkinson's disease. *Mol. Neurobiol.* 45, 173–185. doi: 10.1007/s12035-011-8230-0
- Navailles, S., and De Deurwaerdère, P. (2012b). Imbalanced dopaminergic transmission mediated by serotonergic neurons in L-DOPA-induced dyskinesia. *Parkinsons Dis.* 2012:323686. doi: 10.1155/2012/323686
- Navailles, S., Di Giovanni, G., and De Deurwaerdère, P. (2015). The 5-HT4 agonist prucalopride stimulates L-DOPA-induced dopamine release in restricted brain regions of the hemiparkinsonian rat *in vivo*. *CNS Neurosci. Ther.* 21, 745–747. doi: 10.1111/cns.12436
- Navailles, S., Milan, L., Khalki, H., Di Giovanni, G., Lagière, M., and De Deurwaerdère, P. (2014). Noradrenergic terminals regulate L-DOPA-derived dopamine extracellular levels in a region-dependent manner in Parkinsonian rats. *CNS Neurosci. Ther.* 20, 671–678. doi: 10.1111/cns.12275
- Nevalainen, N., Af Bjerkén, S., Lundblad, M., Gerhardt, G. A., and Strömberg, I. (2011). Dopamine release from serotonergic nerve fibers is reduced in L-DOPA-induced dyskinesia. *J. Neurochem.* 118, 12–23. doi: 10.1111/j.1471-4159.2011.07292.x
- Ng, K. Y., Chase, T. N., Colburn, R. W., and Kopin, I. J. (1970). L-Dopa-induced release of cerebral monoamines. *Science* 170, 76–77. doi: 10.1126/science.170.3953.76
- Politis, M., Wu, K., Loane, C., Brooks, D. J., Kiferle, L., Turkheimer, F. E., et al. (2014). Serotonergic mechanisms responsible for levodopa-induced dyskinesias in Parkinson's disease patients. *J. Clin. Invest.* 124, 1340–1349. doi: 10.1172/JCI71640
- Porras, G., De Deurwaerdere, P., Li, Q., Marti, M., Morgenstern, R., Sohr, R., et al. (2014). L-dopa-induced dyskinesia: beyond an excessive dopamine tone in the striatum. *Sci. Rep.* 4:3730. doi: 10.1038/srep03730
- Prinz, A., Selesnew, L.-M., Liss, B., Roeper, J., and Carlsson, T. (2013). Increased excitability in serotonin neurons in the dorsal raphe nucleus in the 6-OHDA mouse model of Parkinson's disease. *Exp. Neurol.* 248, 236–245. doi: 10.1016/j.expneurol.2013.06.015
- Reed, M. C., Nijhout, H. F., and Best, J. A. (2012). Mathematical insights into the effects of levodopa. *Front. Integr. Neurosci.* 6:21. doi: 10.3389/fnint.2012.00021
- Richard, I. H., McDermott, M. P., Kurlan, R., Lyness, J. M., Como, P. G., Pearson, N., et al. (2012). A randomized, double-blind, placebo-controlled trial of antidepressants in Parkinson disease. *Neurology* 78, 1229–1236. doi: 10.1212/WNL.0b013e3182516244
- Rocha, F. L., Murad, M. G. R., Stumpf, B. P., Hara, C., and Fuzikawa, C. (2013). Antidepressants for depression in Parkinson's disease: systematic review and meta-analysis. *J. Psychopharmacol.* 27, 417–423. doi: 10.1177/0269881113478282
- Rylander, D., Parent, M., O'Sullivan, S. S., Dovero, S., Lees, A. J., Bezard, E., et al. (2010). Maladaptive plasticity of serotonin axon terminals in levodopa-induced dyskinesia. *Ann. Neurol.* 68, 619–628. doi: 10.1002/ana.22097
- Scholtissen, B., Verhey, F. R. J., Adam, J. J., Weber, W., and Leentjens, A. F. G. (2006a). Challenging the serotonergic system in Parkinson disease patients: effects on cognition, mood, and motor performance. *Clin. Neuropharmacol.* 29, 276–285. doi: 10.1097/01.wnf.0000229013.95927.c7
- Scholtissen, B., Verhey, F. R. J., Steinbusch, H. W. M., and Leentjens, A. F. G. (2006b). Serotonergic mechanisms in Parkinson's disease: opposing results from preclinical and clinical data. *J. Neural. Transm. (Vienna)* 113, 59–73. doi: 10.1007/s00702-005-0368-3
- Skapinakis, P., Bakola, E., Salanti, G., Lewis, G., Kyritsis, A. P., and Mavreas, V. (2010). Efficacy and acceptability of selective serotonin reuptake inhibitors for the treatment of depression in Parkinson's disease: a systematic review and meta-analysis of randomized controlled trials. *BMC Neurol.* 10:49. doi: 10.1186/1471-2377-10-49
- Smith, R., Wu, K., Hart, T., Loane, C., Brooks, D. J., Björklund, A., et al. (2015). The role of pallidal serotonergic function in Parkinson's disease dyskinesias: a positron emission tomography study. *Neurobiol. Aging* 36, 1736–1742. doi: 10.1016/j.neurobiolaging.2014.12.037
- Stansley, B. J., and Yamamoto, B. K. (2014). Chronic L-dopa decreases serotonin neurons in a subregion of the dorsal raphe nucleus. *J. Pharmacol. Exp. Ther.* 351, 440–447. doi: 10.1124/jpet.114.218966
- Stansley, B. J., and Yamamoto, B. K. (2015a). Behavioral impairments and serotonin reductions in rats after chronic L-dopa. *Psychopharmacology (Berl)* 232, 3203–3213. doi: 10.1007/s00213-015-3980-4
- Stansley, B. J., and Yamamoto, B. K. (2015b). L-Dopa and brain serotonin system dysfunction. *Toxics* 3, 75–88. doi: 10.3390/toxics3010075
- Svenningsson, P., Rosenblad, C., Af Edholm Arvidsson, K., Wictorin, K., Keywood, C., Shankar, B., et al. (2015). Eltopazine counteracts L-DOPA-induced dyskinesias in Parkinson's disease: a dose-finding study. *Brain* 138, 963–973. doi: 10.1093/brain/awu409
- Tanaka, H., Kannari, K., Maeda, T., Tomiyama, M., Suda, T., and Matsunaga, M. (1999). Role of serotonergic neurons in L-DOPA-derived extracellular dopamine in the striatum of 6-OHDA-lesioned rats. *Neuroreport* 10, 631–634. doi: 10.1097/00001756-199902250-00034
- Wang, S., Zhang, Q. J., Liu, J., Wu, Z. H., Wang, T., Gui, Z. H., et al. (2009). Unilateral lesion of the nigrostriatal pathway induces an increase of neuronal firing of the midbrain raphe nuclei 5-HT neurons and a decrease of their response to 5-HT(1A) receptor stimulation in the rat. *Neuroscience* 159, 850–861. doi: 10.1016/j.neuroscience.2008.12.051
- Yamato, H., Kannari, K., Shen, H., Suda, T., and Matsunaga, M. (2001). Fluoxetine reduces L-DOPA-derived extracellular DA in the 6-OHDA-lesioned rat striatum. *Neuroreport* 12, 1123–1126. doi: 10.1097/00001756-200105080-00015
- Zhang, Q.-J., Gao, R., Liu, J., Liu, Y.-P., and Wang, S. (2007). Changes in the firing activity of serotonergic neurons in the dorsal raphe nucleus in a rat model of Parkinson's disease. *Sheng Li Xue Bao* 59, 183–189.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Migueluez, Benazzouz, Ugedo and De Deurwaerdère. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.